Site Index Estimates

By Site Series:

Report by Region

(2005 Approximation)

Ministry of Forests
Forest Science Program

2005 Approximation Site Index-Site Unit Reports

This document presents an update to the 2003 approximation of estimates of average site index for tree species according to site units of the Biogeoclimatic Ecosystem Classification (BEC) system of British Columbia. The estimates are presented in two site index-site unit table formats: Report by Region and Report by Biogeoclimatic Unit.

In the **Site Index-Site Unit Report by Region**, the tables for biogeoclimatic subzones and variants are presented alphabetically by forest region and biogeoclimatic units. There is one table for each forested biogeoclimatic subzone/variant in the region. In the **Site Index-Site Unit Report by Biogeoclimatic Unit**, the tables are presented alphabetically by biogeoclimatic subzones and variants.

For those site series and species combinations within each BEC subzone/variant that meet the minimum sample size criteria of 7 samples, point estimates of mean site index and their standard error are reported along with their sample size. Otherwise the first approximation site index class estimates are reported.

Note:
1. The following Biogeoclimatic Units have recommended upper elevation limits when applying the SIBEC estimates:

ESSFdc1 – 1850m (Kamloops)
ESSFdc1 – 1800 (Nelson)
ESSFdc2 – 1750m
ESSFdv – 1750m
ESSFdk – 1850m
ESSFmc – 1400m
ESSFmk – 1400m
ESSFmm1 – 1400m
ESSFmw – 1550m
ESSFmv1 – 1300m
ESSFmv2 – 1250m
ESSFmv3 – 1300m
ESSFmv4 – 1250m
ESSFwc2 – 1500m (Prince George)
ESSFwc2 – 1600m (Nelson)
ESSFwc3 – 1450m
ESSFwc4 – 1800m
ESSFwk2 – 1150m
ESSFwm – 1750m
ESSFwv – 1300m
ESSFvc – 1550m
ESSFxc – 1950m

Table of Contents

Report By Region

Region, Biogeoclimatic Unit	Page
Cariboo - ESSFwc3	1
Cariboo - ESSFwk1	2
Cariboo - ESSFxv1	3
Cariboo - ICHdk	4
Cariboo - ICHmk3	5
Cariboo - ICHwk2	6
Cariboo - ICHwk4	7
Cariboo - IDFdk3	8
Cariboo - IDFdk4	9
Cariboo - IDFdw	10
Cariboo - IDFxm	11
Cariboo - IDFxw	12
Cariboo - MSdc2	13
Cariboo - MSdv	14
Cariboo - MSxk	15
Cariboo - MSxv	16
Cariboo - SBPSdc	17
Cariboo - SBPSmk	18
Cariboo - SBPSxc	19
Cariboo - SBSdw1	20
Cariboo - SBSdw2	21
Cariboo - SBSmc1	22
Cariboo - SBSmc2	23
Cariboo - SBSmh	24
Cariboo - SBSmw	25
Cariboo - SBSwk1	26
Kamloops - BGxh1	27
Kamloops - BGxh2	28
Kamloops - BGxw1	29
Kamloops - ESSFdc1	30
Kamloops - ESSFdc2	31
Kamloops - ESSFdv	32
Kamloops - ESSFmw	33
Kamloops - ESSFvc	34
Kamloops - ESSFvv	36
Kamloops - ESSFwc1	37
Kamloops - ESSFwc2	38
Kamloops - ESSFwc4	39
Kamloops - ESSFxc	40
Kamloops - ICHmk1	41
Kamloops - ICHmk2	42
Kamloops - ICHmw2	44
Kamloops - ICHmw3	45
Kamloops - ICHvk1	47
Kamloops - ICHwk1	49
Kamloops - IDFdk1	51
Kamloops - IDFdk2	53
Kamloops - IDFdk3	54
Kamloops - IDFdm1	55
Kamloops - IDFmw1	56
Kamloops - IDFmw2	58
Kamloops - IDFww	60
Kamloops - IDFxh1	61
Kamloops - IDFxh2	63

Kamloops - MSdc 64
Kamloops - MSdm1 65
Kamloops - MSdm2 66
Kamloops - MSxk 67
Kamloops - PPxh1 68
Kamloops - PPxh2 69
Kamloops - SBSmm 70
Nelson - ESSFdc1 71
Nelson - ESSFdk 72
Nelson - ESSFvc 73
Nelson - ESSFwc1 75
Nelson - ESSFwc2 76
Nelson - ESSFwc4 77
Nelson - ESSFwm 78
Nelson - ICHdw 79
Nelson - ICHmk1 81
Nelson - ICHmw1 83
Nelson - ICHmw2 85
Nelson - ICHmw3 87
Nelson - ICHvk1 89
Nelson - ICHwk1 91
Nelson - ICHxw 93
Nelson - IDFdm1 95
Nelson - IDFdm2 97
Nelson - IDFxh1 99
Nelson - MSdk 100
Nelson - MSdm1 101
Nelson - PPdh1 103
Nelson - PPdh2 104
Prince George - BWBSdk1 105
Prince George - BWBSdk2 106
Prince George - BWBSmw1 107
Prince George - BWBSmw2 108
Prince George - BWBSwk1 109
Prince George - BWBSwk2 110
Prince George - BWBSwk3 111
Prince George - ESSFmm1 112
Prince George - ESSFmv1 113
Prince George - ESSFmv2 114
Prince George - ESSFmv3 115
Prince George - ESSFmv4 116
Prince George - ESSFwc2 117
Prince George - ESSFwc3 118
Prince George - ESSFwk1 119
Prince George - ESSFwk2 120
Prince George - ICHmm 121
Prince George - ICHvk2 122
Prince George - ICHwk1 124
Prince George - ICHwk3 126
Prince George - ICHwk4 128
Prince George - SBPSmc 130
Prince George - SBSdh 131
Prince George - SBSdh1 132
Prince George - SBSdk 134
Prince George - SBSdw1 135
Prince George - SBSdw2 136
Prince George - SBSdw3 137
Prince George - SBSmc2 138
Prince George - SBSmc3 140

Prince George - SBSmh 141
Prince George - SBSmk1 142
Prince George - SBSmk2 143
Prince George - SBSmw 145
Prince George - SBSvk 146
Prince George - SBSwk1 147
Prince George - SBSwk2 148
Prince George - SBSwk3 149
Prince Rupert - BWBSdk1 150
Prince Rupert - BWBSdk2 151
Prince Rupert - CWHvh2 152
Prince Rupert - CWHvm1 153
Prince Rupert - CWHvm2 155
Prince Rupert - CWHwm 157
Prince Rupert - CWHws1 159
Prince Rupert - CWHws2 161
Prince Rupert - ESSFmc 162
Prince Rupert - ESSFmk 163
Prince Rupert - ESSFmv3 164
Prince Rupert - ESSFwv 166
Prince Rupert - ICHmc1 167
Prince Rupert - ICHmc1a 168
Prince Rupert - ICHmc2 169
Prince Rupert - ICHvc 170
Prince Rupert - ICHwc 171
Prince Rupert - MHmm1 172
Prince Rupert - MHmm2 173
Prince Rupert - MHwh1 174
Prince Rupert - SBPSmc 175
Prince Rupert - SBSdk 176
Prince Rupert - SBSmc2 177
Prince Rupert - SBSwk3 178
Vancouver - CDFmm 179
Vancouver - CWHdm 180
Vancouver - CWHds1 181
Vancouver - CWHds2 183
Vancouver - CWHmm1 185
Vancouver - CWHmm2 186
Vancouver - CWHms1 188
Vancouver - CWHms2 190
Vancouver - CWHvh1 192
Vancouver - CWHvh2 194
Vancouver - CWHvm1 196
Vancouver - CWHvm2 198
Vancouver - CWHwh1 200
Vancouver - CWHwh2 202
Vancouver - CWHws2 203
Vancouver - CWHxm 204
Vancouver - ESSFmw 206
Vancouver - IDFww 208
Vancouver - MHmm1 209
Vancouver - MHmm2 211
Vancouver - MHwh 212

[redacted] ESSF wc3

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Oak fern		15.0			12.0			15.0	
02	Bl - Rhododendron - Queen's cup		9.0			9.0			9.0	
03	Bl - Globeflower - Horsetail		15.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Oak fern - Brachythecium		15.0			18.0			15.0	
02	Bl - Huckleberry - Feathermoss		9.0			12.0			9.0	
03	Bl - Oak fern - Knight's plume		12.0			15.0			12.0	
04	Bl - Twinberry - Lady fern		15.0			18.0			15.0	
05	Bl - Devil's club - Lady fern		15.0			18.0			15.0	
06	Bl - Horsetail - Sphagnum		9.0						9.0	
07	Bl - Lady fern - Horsetail		12.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Arnica - Cladonia		12.0			15.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwSxw - Falsebox - Wintergreen		24.0			24.0			21.0	
02	CwSxw - Soopolallie		15.0			15.0				
03	CwSxw - Falsebox - Soopolallie					18.0				
04	CwSxw - Falsebox - Feathermoss		21.0			21.0			21.0	
05	CwSxw - Thimbleberry		24.0			24.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwSxw - Falsebox - Knight's plume		24.0			24.0			21.0	
02	FdCw - Wavy-leaved moss		15.0			18.0				
03	CwSxw - Soopolallie		18.0			21.0				
04	CwSxw - Oak fern - Cat's-tail moss		24.0			24.0			24.0	
05	SxwCw - Oak fern					21.0			21.0	
06	CwHw - Devil's club - Lady fern		21.0			21.0			24.0	
07	CwSxw - Devil's club - Horsetail					21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0			21.0	
02	HwCw - Cladonia		15.0			15.0				
03	CwFd - Juniper - Falsebox		15.0			15.0				
04	HwCw - Step moss		21.0			21.0			15.0	
05	SxwCw - Oak fern					21.0			21.0	
06	Sxw - Twinberry - Oak fern					21.0			21.0	
07	CwHw - Devil's club - Lady fern		24.0			21.0			24.0	
08	CwSxw - Skunk cabbage								21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0			21.0	
02	HwCw - Cladonia		15.0			15.0				
03	CwSxw - Soopolallie		15.0			15.0				
04	CwSxw - Velvet-leaved blueberry		18.0			21.0			18.0	
05	HwCw - Step moss		21.0			21.0			18.0	
06	Sxw - Twinberry - Oak fern					21.0			21.0	
07	CwHw - Devil's club - Lady fern		24.0			21.0			24.0	
08	Sxw - Devil's club - Lady fern								18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Feathermoss		15.0			18.0				
02	Fd - Juniper - Kinnikinnick		12.0			12.0				
03	Fd - Juniper - Peltigera		12.0			15.0				
04	Fd - Bluebunch wheatgrass - Needlegrass		12.0							
05	Fd - Feathermoss - Step moss		15.0			15.0				
06	Fd - Pinegrass - Aster		18.0			18.0				
07	SxwFd - Prickly rose - Sedge		18.0			18.0			18.0	
08	SxwFd - Prickly rose - Sarsaparilla		18.0			18.0			18.0	
09	Sxw - Horsetail - Glow moss					18.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Feathermoss		12.0			12.0				
02	Fd - Juniper - Peltigera		9.0			9.0				
03	Fd - Juniper - Saskatoon		9.0							
04	Fd - Juniper - Pasture sage		9.0							
05	Fd - Bluebunch wheatgrass - Pinegrass		9.0			9.0				
06	Pl - Kinnikinnick - Cladonia					9.0				
07	Fd - Feathermoss - Step moss		12.0			12.0				
08	Sxw - Scrub birch - Feathermoss					12.0			9.0	
09	Sxw - Feathermoss - Brachythecium		15.0			15.0			15.0	
10	Sxw - Horsetail - Glow moss								15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Fd - Prickly rose - Pinegrass		15.0			18.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Fd - Pinegrass - Feathermoss		15.0							
02	Fd - Bluebunch wheatgrass - Penstemon		12.0							
03	Fd - Juniper - Cladonia		12.0			12.0				
04	Fd - Bluebunch wheatgrass - Pasture sage		12.0							
05	Fd - Feathermoss - Step moss		15.0							
06	Fd - Ricegrass - Feathermoss		15.0							
07	Fd - Prickly rose - Sarsaparilla		15.0							
08	Sxw - Snowberry - Prickly rose		18.0						18 0	
09	Sxw - Horsetail								18 0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Fd - Juniper - Bluebunch wheatgrass		15.0			12.0				
02	FdPy - Bluebunch wheatgrass - Pinegrass		6.0			6.0				
03	FdPy - Western snowberry - Bluebunch whe		9.0			9.0				
04	FdPy - Bluebunch wheatgrass - Balsamroot		9.0			9.0				
05	Fd - Feathermoss		15.0							
06	Sxw - Waterbirch		21.0						18.0	
07	Sxw - Prickly rose - Coltsfoot		21.0						21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Wintergreen - Feathermoss		15.0			18.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] MS

Site Series(*)	Name	Bl			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Soopolallie - Twinflower		15.0			18.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Pinegrass - Lupine		12.0			15.0		20	15.7	0.6	12	14.8	0.6
02	Fd - Juniper - Grouseberry		9.0			12.0		10	12.4	0.9		12.0	
05	FdPl - Pinegrass - Arnica		12.0			15.0		8	14.9	0.6		12.0	
06	Pl - Falsebox - Lupine		15.0			18.0		27	16.0	0.5	7	13.0	0.8
07	Sxw - Trapper's tea - Grouseberry		12.0					14	17.0	0.9		12.0	
08	Sxw - Gooseberry - Grouseberry		15.0						18.0			15.0	
09	Sxw - Horsetail - Leafy moss		12.0						15.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Sx		
		#	SI	SE	#	SI	SE
01	Pl - Grouseberry - Feathermoss		18.0			12.0	
02	Pl - Fescue - Stereocaulon		12.0				
03	Pl - Kinnikinnick - Cladonia		12.0				
04	Pl - Grouseberry - Kinnikinnick		15.0			9.0	
05	Pl - Trapper's tea - Crowberry		18.0			12.0	
06	Sxw - Crowberry - Knight's plume		18.0			15.0	
07	Sxw - Crowberry - Glow moss		12.0			12.0	
08	Sxw - Horsetail - Crowberry		12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Sx		
		#	SI	SE	#	SI	SE
01	Pl - Juniper - Feathermoss		18.0			15.0	
02	Pl - Kinnikinnick - Cladonia		12.0				
03	Pl - Kinnikinnick - Feathermoss		15.0				
04	PlSb - Feathermoss		15.0				
05	Sxw - Scrub birch - Feathermoss		15.0			15.0	
06	Sxw - Horsetail - Meadowrue		21.0			21.0	
07	Sb - Scrub birch - Sedge		9.0			12.0	
08	Sxw - Horsetail - Glow moss		15.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SBPSmk

Site Series(*)	Name	Pl			Sx			Fd		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Pinegrass - Arnica		18.0			15.0				
02	Pl - Cladonia - Haircap moss		15.0			9.0				
03	Fd - Pinegrass - Aster		18.0						15.0	
04	Pl - Pinegrass - Feathermoss		18.0			15.0				
05	SxwFd - Step moss		21.0			15.0			18.0	
06	Sxw - Twinberry		21.0			21.0				
07	Sxw - Horsetail - Glow moss		18.0			21.0				
08	Sb - Scrub birch - Sedge		12.0			12.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Fd			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Kinnikinnick - Feathermoss		15.0							
02	Pl - Kinnikinnick - Cladonia		12.0			12 0				
03	Sxw - Scrub birch - Fen moss		15.0						12 0	
04	Sxw - Scrub birch - Feathermoss		15.0						15 0	
05	Sxw - Horsetail - Glow moss		15.0						15 0	
06	Sxw - Horsetail - Meadowrue		21.0						18 0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass		21.0			24.0			21.0	
02	FdPl - Cladonia		15.0			15.0				
03	Fd - Saskatoon - Pinegrass		15.0			15.0				
04	Pl - Pinegrass - Feathermoss		15.0			18.0			15.0	
05	SxwFd - Ricegrass		21.0			21.0			18.0	
06	SxwFd - Thimbleberry		21.0			24.0			21.0	
07	Sxw - Twinberry - Coltsfoot		21.0			21.0			21.0	
08	Sxw - Twinberry - Oak fern		21.0			24.0			24.0	
09	Sxw - Horsetail - Glow moss					21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass		18.0			21.0			18.0	
02	FdPl - Cladonia		15.0			15.0				
03	Pl - Kinnikinnick - Wavy-leaved moss					15.0				
04	Fd - Pinegrass - Aster		15.0			15.0				
05	SxwFd - Cat's-tail moss		18.0			21.0			18.0	
06	Pl - Pinegrass - Feathermoss		18.0			21.0			18.0	
07	PlSb - Feathermoss					18.0			15.0	
08	Sxw - Twinberry		18.0			21.0			18.0	
09	Sxw - Devil s club - Knight's plume		21.0			21.0			21.0	
10	Sxw - Horsetail					15.0			18.0	
11	Sb - Soft-leaved sedge - Sphagnum					9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry		15.0			18.0			18.0			18.0	
02	Pl - Cladonia - Haircap moss		12.0						15.0			12.0	
03	Fd - Pinegrass - Aster					18.0			15.0				
04	Sxw - Huckleberry - Labrador tea		15.0						18.0			15.0	
05	Sxw - Spirea - Glow moss		15.0						15.0			15.0	
06	Sxw - Oak fern		18.0			18.0			21.0			21.0	
07	Sxw - Devil's club - Step moss		18.0			21.0			21.0			21.0	
08	Sxw - Horsetail - Glow moss		18.0						15.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry	31	15.4	0.5	97	17.9	0.3	48	18.7	0.5
02	Pl - Huckleberry - Cladonia		12.0		27	16.4	0.5	7	12.3	0.8
04	Sxw - Huckleberry - Dwarf blueberry					18.0			18.0	
06	Sxw - Oak fern	17	19.1	0.7	10	19.9	0.5	26	19.3	0.5
08	Sxw - Twinberry - Oak fern					21.0			21.0	
11	Sxw - Horsetail - Glow moss					18.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSmh

Site Series(*)	Name	Fd			Sx			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Hazelnut		21.0			18.0				
02	FdPl - Cladonia		18.0							
03	FdPl - Velvet-leaved blueberry - Cladonia		18.0			15.0			21.0	
04	Fd - Douglas maple - Step moss		21.0			15.0				
05	SxwFd - Feathermoss		21.0			18.0				
06	SxwFd - Coltsfoot		21.0			21.0				
07	SxwEp - Devil's club		24.0			24.0				
08	Sxw - Ostrich fern		24.0			24.0				
09	Sxw - Horsetail - Glow moss					21.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSxw

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Falsebox		15.0		8	22.6	0.6	15	22.4	0.6	22	20.0	0.4
02	FdBl - Huckleberry					15.0			15.0			15.0	
03	Pl - Huckleberry - Velvet-leaved blueberry					18.0		8	17.1	1.2		9.0	
04	SxwFd - Knight's plume					18.0			21.0			15.0	
05	Sxw - Pink spirea							7	18.2	0.7		15.0	
06	Sxw - Oak fern		15.0			21.0			24.0			21.0	
07	Sxw - Twinberry - Oak fern		15.0						24.0			21.0	
08	Sxw - Devil's club		18.0			21.0			24.0			24.0	
09	Sxw - Horsetail		15.0						18.0			18.0	
10	SbSxw - Scrub birch - Sedge								9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSwk1

Site Series(*)	Name	Bl			Fd			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Oak fern		15.0			21.0			21.0		13	21.2	0.8			
02	Pl - Huckleberry - Cladina		12.0			15.0			15.0			12.0				
03	Pl - Huckleberry - Velvet-leaved blueberry		12.0			15.0			18.0			15.0				
04	SxwFd - Knight's plume		15.0			18.0			18.0			15.0				
05	Sxw - Huckleberry - Highbush-cranberry		15.0						21.0			18.0				
06	Sxw - Pink spirea - Oak fern		15.0						21.0			18.0				
07	Sxw - Twinberry - Oak fern		18.0						21.0			18.0				
08	Sxw - Devil's club		18.0						21.0			18.0				
09	Sxw - Horsetail		18.0						21.0			18.0				
11	SbSxw - Scrub birch - Sedge								12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops BGxh1

Site Series(*)	Name	Fd			Py		
		#	SI	SE	#	SI	SE
04	Py - Antelope brush - Red three-awn		9.0			12.0	
05	Py - Sumac		15.0			15.0	
06	PyAct - Nootka rose - Poison ivy		18.0			18.0	
07	Act - Water birch		21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Py			Fd		
		#	SI	SE	#	SI	SE
03	Py - Red three-awn		12.0				
04	Py - Bluebunch wheatgrass		12.0			12.0	
07	Act - Snowberry - Dogwood		21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py		
		#	SI	SE	#	SI	SE
03	Py - Bluebunch wheatgrass		12.0			12.0	
05	Py - Rough fescue - Bluebunch wheatgrass		15.0			15.0	
08	At - Snowberry - Kentucky bluegrass		21.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Grouseberry		12.0			15.0			12.0	
02	PlSe - Pinegrass		9.0			12.0			9.0	
03	Bl - Grouseberry - Cladonia		9.0			12.0			9.0	
04	Bl - Rhododendron - Valerian		12.0			15.0			12.0	
05	Bl - Trapper's tea		12.0			12.0			12.0	
06	Bl - Horsetail - Glow moss		9.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFdc2

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Grouseberry	9	16.4	0.8	9	17.3	0.6	14	16.1	1.0
03	PlSe - Falsebox - Pinegrass		12.0			15.0			12.0	
04	Bl - Grouseberry - Cladonia	7	14.2	1.5		15.0			12.0	
05	Bl - Huckleberry - Feathermoss		12.0			15.0			12.0	
06	Bl - Gooseberry - Oak fern		12.0			15.0		9	18.6	1.4
07	Bl - Rhododendron - Valerian		12.0			15.0			15.0	
08	Bl - Trapper's tea		9.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] ESSFdv

Site Series(*)	Name	Bl			Pl			Se			Pa		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Heron's-bill		12.0			15.0			12.0				
02	Pa - Junegrass		9.0			12.0			9.0			6.0	
03	BlPa - Juniper		9.0			12.0			9.0			9.0	
04	Bl - Huckleberry - Brachythecium		12.0			15.0			12.0			9.0	
05	Bl - Valerian - Arnica		12.0			15.0			15.0				
06	Bl - Horsetail - Glow moss		9.0			9.0			9.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFmw

Site Series(*)	Name	Ba			Bl			Pl			Se			Pa		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	BlBa - Rhododendron		15.0			15.0			15.0			15.0				
02	BlPl - Juniper - Rhacomitrium					9.0			12.0			12.0			9.0	
03	FdBl - Falsebox - Pinegrass								15.0			12.0				
04	Bl - Huckleberry - Falsebox					12.0			15.0			12.0				
05	BlBa - Azalea - Pipecleaner moss		15.0			15.0			18.0			15.0				
06	Bl - Gooseberry - Valerian		15.0			15.0			15.0			15.0				
07	BlBa - Oak fern - Lady fern		18.0			18.0						18.0				
08	Bl - Gooseberry - Horsetail											18.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFmw

Site Series(*)	Name	Fd		
		#	SI	SE
01	BlBa - Rhododendron			
02	BlPl - Juniper - Rhacomitrium			
03	FdBl - Falsebox - Pinegrass		15.0	
04	Bl - Huckleberry - Falsebox		15.0	
05	BlBa - Azalea - Pipecleaner moss			
06	Bl - Gooseberry - Valerian			
07	BlBa - Oak fern - Lady fern			
08	Bl - Gooseberry - Horsetail			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFvc

Site Series(*)	Name	Bl			Hm			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	BlHm - Rhododendron - Oak fern		15.0			12.0			15.0	
02	BlHm - Rhododendron - Leafy liverwort		12.0			9.0			12.0	
03	BlHm - Rhododendron - Pipecleaner moss		15.0			12.0			15.0	
04	BlHm - Devil's club - Lady fern		18.0						18.0	
05	BlHm - Horsetail		18.0						18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Se			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Foamflower		12.0			12.0				
02	Bl - Huckleberry - Mountain liverwort		9.0			9.0			12.0	
03	BlHm - Rhododendron - Leafy liverwort		12.0			9.0				
04	Bl - Valerian - Groundsel		12.0			12.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] ps 00St wc1

Site Series(*)	Name	Bl			Cw			Hw			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Oak fern	18	19.6	0.7		15.0			15.0		18	20.5	0.3	18	18.5	0.8
02	Bl - Falsebox - Grouseberry	29	17.2	0.7							28	17.8	0.6		15.0	
03	Bl - Devil's club - Lady fern	15	19.7	0.5		15.0			15.0			18.0			21.0	
04	Bl - Horsetail - Brachythecium		18.0						15.0			18.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFwc2

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Oak fern		15.0			15.0			15.0	
02	Pl - Huckleberry - Cladonia		9.0			12.0			12.0	
03	Bl - Huckleberry - Arnica		12.0			15.0			12.0	
04	Bl - Rhododendron - Heron's-bill		12.0			15.0			12.0	
05	Bl - Azalea - Feathermoss	11	13.6	0.8		12.0				
06	Bl - Valerian - Oak fern					18.0				
07	Bl - Devil's club - Lady fern		18.0			18.0			21.0	
08	Bl - Horsetail - Sphagnum		15.0			12.0			15.0	
09	Pl - Dwarf blueberry - Sphagnum		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFwc4

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Oak fern	8	16.2	1.2		21.0			18.0	
02	Bl - Rhododendron - Falsebox	11	15.6	0.9	14	15.8	0.6		15.0	
03	Bl - Rhododendron - Woodrush		15.0			18.0			15.0	
04	Bl - Rhododendron - Foamflower		15.0			21.0			18.0	
05	Bl - Rhododendron - Lady fern		18.0						18.0	
06	Bl - Horsetail - Brachythecium		15.0						18.0	
07	Bl - Sedge - Sphagnum		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ESSFxc

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Grouseberry - Valerian	9	15.6	1.1	8	16.1	1.1		12.0	
02	Pl - Juniper - Lupine		9.0			12.0			9.0	
05	Bl - Grouseberry - Cladonia	7	16.2	1.3	11	16.9	1.3		12.0	
06	Bl - Rhododendron - Grouseberry	11	13.0	0.9		15.0		8	15.3	0.8
07	Bl - Gooseberry - Foamflower		15.0			18.0			15.0	
08	Bl - Horsetail - Glow moss		12.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops (CHmk1

Site Series(*)	Name	Bl			Cw			Fd			Lw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwSxw - Falsebox	10	16.5	1.1		15.0		18	20.4	0.8		21.0		7	19.3	0.9
01-YS	CwSxw - Aspen		18.0			15.0			21.0			21.0			21.0	
02	Fd - Penstemon - Pinegrass								15.0						18.0	
03	FdPl - Pinegrass - Twinflower		15.0						18.0			21.0			21.0	
04	FdPl - Sitka alder - Pinegrass		18.0			12.0		11	22.6	0.9		21.0		16	20.5	0.9
05	Sxw - Gooseberry - Sarsaparilla		21.0			18.0			21.0			24.0			21.0	
06	Sxw - Oak fern		21.0			18.0			24.0			24.0			24.0	
07	Sxw - Horsetail		18.0			15.0									21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ICHmk1

Site Series(*)	Name	Sx		
		#	SI	SE
01	CwSxw - Falsebox	12	19.4	1.2
01-YS	CwSxw - Aspen		18.0	
02	Fd - Penstemon - Pinegrass			
03	FdPl - Pinegrass - Twinflower		18.0	
04	FdPl - Sitka alder - Pinegrass		18.0	
05	Sxw - Gooseberry - Sarsaparilla		21.0	
06	Sxw - Oak fern		21.0	
07	Sxw - Horsetail		18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops ICHmk2

Site Series(*)	Name	Bl			Cw			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwSxw - Falsebox - Knight's plume		18.0			15.0			18.0			21.0			18.0	
02	Fd - Juniper - Pinegrass								15.0			18.0				
03	Fd - Falsebox - Pinegrass		15.0						18.0			21.0			18.0	
04	CwSxw - Douglas maple - Fairybells					12.0			18.0			21.0			18.0	
05	CwSxw - Oak fern - Bunchberry		21.0			18.0			21.0			24.0			21.0	
06	Sxw - Horsetail		18.0			15.0						21.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Kamloops (CHmw2 **)

Site Series(*)	Name	Bl			Cw			Fd			Hw			Lw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		18.0			18.0			24.0		9	19.3	1.3		24.0	
01-YS	CwFd - Feathermoss		18.0			18.0			24.0			18.0			24.0	
02	FdCw - Falsebox - Prince's pine		15.0			15.0		17	22.4	0.8	14	19.4	1.0	16	21.1	0.5
03	CwFd - Falsebox		18.0			18.0			21.0			18.0		10	22.3	1.2
04	CwHw - Oak fern - Foamflower		21.0			18.0			24.0			18.0			24.0	
05	CwHw - Devil's club - Lady fern		21.0			27.0			27.0			21.0			27.0	
06	CwHw - Horsetail		18.0									15.0				
07	CwSxw - Skunk cabbage		18.0									15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]ps ...mw2

Site Series(*)	Name	Pl			Pw			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		24.0			24.0			18.0	
01-YS	CwFd - Feathermoss		24.0			24.0			18.0	
02	FdCw - Falsebox - Prince's pine	11	21.2	0.9		24.0			15.0	
03	CwFd - Falsebox		24.0			24.0			18.0	
04	CwHw - Oak fern - Foamflower		24.0			27.0			24.0	
05	CwHw - Devil's club - Lady fern		27.0			27.0			24.0	
06	CwHw - Horsetail								21.0	
07	CwSxw - Skunk cabbage								18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE
01	HwCw - Falsebox - Feathermoss		18.0			15.0			21.0		9	19.4	0.7		21.0	
01-YC	HwCw - Feathermoss		18.0			15.0			21.0			18.0			21.0	
02	Fd - Juniper - Cladina								15.0						18.0	
03	FdPl - Pinegrass - Feathermoss		15.0			15.0			18.0						21.0	
04	CwFd - Soopolallie - Twinflower					15.0			21.0						21.0	
05	CwFd - Falsebox		18.0			15.0			21.0			18.0			21.0	
06	CwHw - Oak fern		21.0			18.0			24.0			18.0			24.0	
07	CwHw - Devil's club - Lady fern		21.0			18.0			27.0			18.0				
08	CwSxw - Skunk cabbage		15.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, Sl = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pw			Sx			Lw		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		24.0			18.0				
01-YC	HwCw - Feathermoss		24.0			18.0				
02	Fd - Juniper - Cladina									
03	FdPl - Pinegrass - Feathermoss					15.0				
04	CwFd - Soopolallie - Twinflower		21.0			18.0			21.0	
05	CwFd - Falsebox		21.0			18.0				
06	CwHw - Oak fern		24.0			21.0				
07	CwHw - Devil's club - Lady fern		27.0			24.0				
08	CwSxw - Skunk cabbage					18.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Devil's club - Lady fern		21.0			21.0			24.0			18.0			24.0	
02	HwCw - Falsebox - Feathermoss		18.0						24.0			18.0			24.0	
03	CwHw - Oak fern - Foamflower		18.0			18.0			24.0			18.0			24.0	
04	CwHw - Oak fern - Spiny wood fern		21.0			18.0			27.0			18.0			27.0	
05	CwSxw - Devil's club - Horsetail		18.0			15.0						15.0			24.0	
06	CwSxw - Skunk cabbage		15.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Sx		
		#	SI	SE
01	CwHw - Devil's club - Lady fern		24.0	
02	HwCw - Falsebox - Feathermoss		21.0	
03	CwHw - Oak fern - Foamflower		21.0	
04	CwHw - Oak fern - Spiny wood fern		24.0	
05	CwSxw - Devil's club - Horsetail		21.0	
06	CwSxw - Skunk cabbage		18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		21.0		10	19.0	1.1		24.0		15	20.3	0.7		24.0	
02	PlHw - Velvet-leaved blueberry		15.0			15.0						15.0			18.0	
03	HwCw - Step moss		18.0			15.0			21.0			15.0			21.0	
04	HwCw - Falsebox - Feathermoss		18.0			18.0		8	23.7	1.7		18.0			21.0	
05	CwHw - Devil's club - Lady fern		21.0			21.0			24.0			18.0				
06	CwSxw - Devil's club - Horsetail		18.0			15.0						15.0				
07	CwSxw - Skunk cabbage		18.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pw			Sx		
		#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0	
02	PlHw - Velvet-leaved blueberry						
03	HwCw - Step moss		24.0			18.0	
04	HwCw - Falsebox - Feathermoss		24.0			21.0	
05	CwHw - Devil's club - Lady fern		27.0			24.0	
06	CwSxw - Devil's club - Horsetail					21.0	
07	CwSxw - Skunk cabbage					18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx			Py			Bl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Feathermoss	9	16.3	0.9	7	18.5	0.9		15.0							
02	Fd - Snowberry - Bluebunch wheatgrass		12.0			15.0						12.0				
03	Fd - Juniper - Pinegrass		15.0			15.0						12.0				
04	Fd - Pinegrass - Feathermoss	11	18.0	0.7	11	17.1	0.8					15.0				
05	SxwFd - Gooseberry - Feathermoss	8	19.9	0.7		21.0		12	20.2	0.8					18.0	
06	Sxw - Horsetail		21.0			21.0			21.0						18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx			Py			Cw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Feathermoss	31	18.2	0.5	21	18.4	0.8		15.0							
02	FdPy - Bluebunch wheatgrass - Pinegrass		12.0									12.0				
03	FdPy - Pinegrass	12	16.3	0.9		18.0						12.0				
04	Fd - Feathermoss	7	18.7	2.0	8	18.2	1.1		15.0			15.0				
05	SxwFd - Dogwood - Gooseberry	7	21.3	1.3		21.0		11	20.4	2.1						
06	Sxw - Horsetail		21.0			21.0			21.0							
07	CwSxw - Twinberry - Soft-leaved sedge					21.0			18.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Feathermoss		15.0			18.0				
02	Fd - Juniper - Kinnikinnick		12.0			12.0				
03	Fd - Juniper - Peltigera		12.0			15.0				
04	Fd - Bluebunch wheatgrass - Needlegrass		12.0							
05	Fd - Feathermoss - Step moss					15.0				
06	Fd - Pinegrass - Aster		15.0			15.0				
07	SxwFd - Prickly rose - Sedge		18.0			18.0			18.0	
08	SxwFd - Prickly rose - Sarsaparilla		18.0			18.0			18.0	
09	Sxw - Horsetail - Glow moss					18.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Lw			Pl			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Twinflower	11	18.3	1.0		18.0			18.0			18.0			15.0	
03	FdPy - Bluebunch wheatgrass - Pinegrass		12.0						12.0			15.0				
04	Fd - Pinegrass - Kinnikinnick		15.0			15.0			15.0			18.0				
05	FdLw - Spruce - Pinegrass		21.0			21.0			21.0			21.0			18.0	
06	SxwFd - Dogwood - Gooseberry		24.0			24.0			24.0						24.0	
07	Sxw - Horsetail		21.0			21.0			21.0						21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	BI		
		#	SI	SE
01	FdPl - Pinegrass - Twinflower			
03	FdPy - Bluebunch wheatgrass - Pinegrass			
04	Fd - Pinegrass - Kinnikinnick			
05	FdLw - Spruce - Pinegrass		18.0	
06	SxwFd - Dogwood - Gooseberry		21.0	
07	Sxw - Horsetail		18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Fd			Lw			Pl			Py		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdCw - Falsebox - Prince's pine		15.0		19	20.6	1.0		24.0		9	21.5	0.8		21.0	
01-YC	CwFd - Feathermoss		15.0			21.0			24.0			21.0			21.0	
02	FdPy - Snowberry - Bluebunch wheatgrass					15.0									15.0	
03	Fd - Penstemon - Pinegrass					18.0						15.0			18.0	
04	Fd - Pinegrass - Feathermoss		15.0		11	18.1	1.3		21.0			18.0			18.0	
05	CwFd - Dogwood		18.0			21.0			24.0			21.0				
06	Cw - Devil's club - Foamflower		21.0			24.0			27.0			21.0				

continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Sx		
		#	SI	SE
01	FdCw - Falsebox - Prince's pine		18.0	
01-YC	CwFd - Feathermoss		18.0	
02	FdPy - Snowberry - Bluebunch wheatgrass			
03	Fd - Penstemon - Pinegrass			
04	Fd - Pinegrass - Feathermoss			
05	CwFd - Dogwood		21.0	
06	Cw - Devil's club - Foamflower		24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Cw			Fd			Pl			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdCw - Falsebox - Prince's pine		15.0		7	23.1	0.9		21.0			18.0			18.0	
02	Fd - Snowberry - Bluebunch wheatgrass					15.0			18.0			15.0				
03	Fd - Pinegrass - Feathermoss					18.0			21.0			15.0				
04	CwSxw - Oak fern		21.0			21.0			24.0						21.0	
05	Dogwood - Sedge		15.0						18.0						18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Fd			Pl			Py			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdCw - Hazelnut		15.0			24.0			24.0			24.0				
02	FdPl - Peltigera					12.0			12.0							
03	Fd - Falsebox - Feathermoss					18.0			18.0			18.0				
04	Fd - Douglas maple - Fairybells		15.0			24.0						24.0				
05	CwFd - Vine maple		24.0			27.0									27.0	
06	Cw - Devil's club - Lady fern		27.0			36.0									36.0	
07	CwSxw - Skunk cabbage		15.0												21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bg		
		#	SI	SE
01	FdCw - Hazelnut			
02	FdPl - Peltigera			
03	Fd - Falsebox - Feathermoss			
04	Fd - Douglas maple - Fairybells			
05	CwFd - Vine maple			
06	Cw - Devil's club - Lady fern		36.0	
07	CwSxw - Skunk cabbage			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	FdPy - Pinegrass	13	16.3	1.1	8	16.8	1.5			
02	FdPy - Bluebunch wheatgrass - Balsamroot		12.0			12.0				
03	FdPy - Bluebunch wheatgrass - Pinegrass		12.0			15.0				
04	FdPy - Snowbrush - Pinegrass		15.0			15.0				
05	FdPy - Pinegrass - Idaho fescue		15.0			15.0				
06	FdPy - Spirea - Feathermoss		18.0			18.0				
07	FdPy - Snowberry - Spirea		18.0			18.0				
08	SxwFd - Douglas maple - Dogwood		21.0			21.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	FdPy - Pinegrass - Feathermoss		15.0			15.0				
02	FdPy - Bluebunch wheatgrass - Rough fesc		12.0			12.0				
03	FdPy - Bluebunch wheatgrass - Balsamroot		12.0			12.0				
04	FdPy - Bluebunch wheatgrass - Pinegrass		15.0			15.0				
05	FdPy - Pinegrass		15.0			15.0				
06	Fd - Feathermoss		15.0			15.0				
07	CwFd - Dogwood		21.0			21.0			18.0	
08	Sxw - Horsetail		21.0						21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Wintergreen - Feathermoss		15.0			15.0			18.0			15.0	
02	FdPl - Juniper		9.0			12.0			12.0			9.0	
03	Pl - Spirea - Pinegrass		12.0			15.0			15.0			12.0	
04	Sxw - Gooseberry		18.0			18.0			21.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Lw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Falsebox - Feathermoss	10	19.3	1.0		18.0			21.0		22	19.2	0.6	14	17.5	1.2
02	Fd - Penstemon - Pinegrass					15.0			15.0			15.0				
03	Pl - Grouseberry - Cladonia		15.0			18.0			18.0		7	13.3	1.7		15.0	
04	Pl - Pinegrass - Kinnikinnick		15.0			18.0			21.0		11	17.6	1.2		15.0	
05	Sxw - Trapper's tea - Grouseberry		18.0			18.0					8	19.7	0.6	7	16.3	0.4
06	Sxw - Gooseberry		18.0			18.0						21.0			18.0	
07	Sxw - Trapper's tea - Horsetail		15.0									18.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Falsebox - Feathermoss	19	17.0	0.7	8	18.9	1.1	27	19.1	0.6	21	16.9	0.7
03	Pl - Juniper - Grouseberry		15.0			18.0		12	15.2	0.9		15.0	
04	Pl - Grouseberry - Pinegrass	7	18.3	1.2	11	19.3	0.4	28	17.2	0.4		15.0	
05	Sxw - Gooseberry - Grouseberry	9	19.3	1.7		18.0		16	20.9	0.5	14	20.6	1.2
06	Sxw - Gooseberry - Devil's club		15.0			18.0			18.0			15.0	
07	Sxw - Horsetail - Leafy moss		12.0						15.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Pinegrass - Lupine		12.0			15.0		20	15.7	0.6	12	14.8	0.6
02	Fd - Juniper - Grouseberry		9.0			12.0		10	12.4	0.9		12.0	
05	FdPl - Pinegrass - Arnica		12.0			15.0		8	14.9	0.6		12.0	
06	Pl - Falsebox - Lupine		15.0			18.0		27	16.0	0.5	7	13.0	0.8
07	Sxw - Trapper's tea - Grouseberry		12.0					14	17.0	0.9		12.0	
08	Sxw - Gooseberry - Grouseberry		15.0						18.0			15.0	
09	Sxw - Horsetail - Leafy moss		12.0						15.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Fd			Py		
		#	SI	SE	#	SI	SE
01	Py - Bluebunch wheatgrass - Idaho fescue		12.0			15.0	
02	Py - Red three-awn					18.0	
04	Py - Bluebunch wheatgrass - Cheatgrass		12.0			15.0	
05	Py - Bluebunch wheatgrass - Rough fescue		18.0			15.0	
06	FdPy - Snowberry - Pinegrass		21.0			18.0	
07	FdPy - Snowberry - Spirea		21.0			18.0	
08	Fd - Water birch - Douglas maple		24.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py		
		#	SI	SE	#	SI	SE
01	Py - Bluebunch wheatgrass - Fescue		12.0			12.0	
02	FdPy - Bluebunch wheatgrass - Selaginella		9.0			9.0	
03	Py - Bluebunch wheatgrass		12.0			12.0	
04	Py - Big sage - Bluebunch wheatgrass		12.0			12.0	
06	FdPy - Snowberry - Saskatoon		18.0			15.0	
07	Act - Water birch		21.0			18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Pl			Sx			Cw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Falsebox - Knight's plume		18.0			18.0			21.0			18.0				
02	Pl - Huckleberry - Cladonia		12.0			15.0			15.0			12.0				
03	Pl - Douglas-fir - Juniper		12.0			15.0			15.0			12.0				
04	Pl - Soopolallie - Pinegrass		15.0			18.0			18.0			15.0				
05	Sxw - Soopolallie - Falsebox		15.0			18.0			18.0			15.0				
06	Sxw - Huckleberry - Falsebox		15.0			18.0			18.0			15.0				
07	Sxw - Oak fern		18.0						21.0			21.0			12.0	
08	Sxw - Horsetail		18.0						18.0			18.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Grouseberry		12.0			15.0			12.0	
02	PlSe - Pinegrass		9.0			12.0			9.0	
03	Bl - Grouseberry - Cladonia		9.0			12.0			9.0	
04	Bl - Rhododendron - Valerian		12.0			15.0			12.0	
05	Bl - Trapper's tea		12.0			12.0			12.0	
06	Bl - Horsetail - Glow moss		9.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Lw			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Foamflower		15.0			18.0			18.0		11	19.8	0.5		15.0	
02	Fd - Douglas maple - Soopolallie		12.0			15.0			15.0			15.0			12.0	
03	Bl - Azalea - Grouseberry	16	14.3	1.0							15	16.3	0.7	14	14.7	0.9
04	Bl - Azalea - Soopolallie	20	14.4	0.8	10	17.3	0.7		18.0		17	17.8	0.8	21	14.3	0.8
05	Bl - Azalea - Step moss		12.0									18.0			15.0	
06	Bl - Azalea - Horsetail		12.0									15.0			15.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pa		
		#	SI	SE
01	Bl - Azalea - Foamflower			
02	Fd - Douglas maple - Soopolallie			
03	Bl - Azalea - Grouseberry			
04	Bl - Azalea - Soopolallie		15.0	
05	Bl - Azalea - Step moss			
06	Bl - Azalea - Horsetail			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Hm			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	BlHm - Rhododendron - Oak fern		15.0			12.0			15.0	
02	BlHm - Rhododendron - Leafy liverwort		12.0			9.0			12.0	
03	BlHm - Rhododendron - Pipecleaner moss		15.0			12.0			15.0	
04	BlHm - Devil's club - Lady fern		18.0						18.0	
05	BlHm - Horsetail		18.0						18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Cw			Hw			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Oak fern	18	19.6	0.7		15.0			15.0		18	20.5	0.3	18	18.5	0.8
02	Bl - Falsebox - Grouseberry	29	17.2	0.7							28	17.8	0.6		15.0	
03	Bl - Devil's club - Lady fern	15	19.7	0.5		15.0			15.0			18.0			21.0	
04	Bl - Horsetail - Brachythecium		18.0						15.0			18.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Oak fern		15.0			15.0			15.0	
02	Pl - Huckleberry - Cladonia		9.0			12.0			12.0	
03	Bl - Huckleberry - Arnica		12.0			15.0			12.0	
04	Bl - Rhododendron - Heron's-bill		12.0			15.0			12.0	
05	Bl - Azalea - Feathermoss	11	13.6	0.8		15.0			12.0	
06	Bl - Valerian - Oak fern		15.0			18.0			15.0	
07	Bl - Devil's club - Lady fern		18.0			18.0			21.0	
08	Bl - Horsetail - Sphagnum		15.0			12.0			15.0	
09	Pl - Dwarf blueberry - Sphagnum		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Oak fern	8	16.2	1.2		21.0			18.0	
02	Bl - Rhododendron - Falsebox	11	15.6	0.9	14	15.8	0.6		15.0	
03	Bl - Rhododendron - Woodrush		15.0			18.0			15.0	
04	Bl - Rhododendron - Foamflower		15.0			21.0			18.0	
05	Bl - Rhododendron - Lady fern		18.0						18.0	
06	Bl - Horsetail - Brachythecium		15.0						18.0	
07	Bl - Sedge - Sphagnum		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Se			Fd			Hw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Arnica		12.0			18.0			15.0							
02	Bl - Rhododendron - Azalea		12.0			15.0			15.0							
03	BlHw - Rhododendron - Azalea		12.0			18.0			15.0			18.0			15.0	
04	Bl - Azalea - Queen's cup		15.0			18.0			18.0							

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Lw			Pw		
		#	SI	SE	#	SI	SE
01	Bl - Azalea - Arnica						
02	Bl - Rhododendron - Azalea						
03	BlHw - Rhododendron - Azalea		18.0			18.0	
04	Bl - Azalea - Queen's cup						

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bg			Bl			Cw			Fd			Hw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01a	CwFd - Falsebox, SX-SM					18.0			15.0			21.0			18.0	
01b	CwFd - Falsebox, M-SHG		21.0			24.0			18.0			24.0			18.0	
02	FdPy - Oregon grape - Parsley fern								18.0			15.0			18.0	
03	CwHw - White pine - Devil's club		21.0			21.0			18.0			27.0			21.0	
04	CwHw - Devil's club - Lady fern		21.0			21.0			21.0			27.0			18.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Lw			Pl			Pw			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01a	CwFd - Falsebox, SX-SM		21.0			21.0			21.0			21.0				
01b	CwFd - Falsebox, M-SHG		24.0			24.0			24.0			24.0			21.0	
02	FdPy - Oregon grape - Parsley fern		21.0			21.0						21.0				
03	CwHw - White pine - Devil's club		27.0			24.0			24.0						24.0	
04	CwHw - Devil's club - Lady fern		27.0			24.0			27.0						24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Lw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwSxw - Falsebox	10	16.5	1.1		15.0		18	20.4	0.8		21.0		7	19.3	0.9
02	Fd - Penstemon - Pinegrass								15.0						18.0	
03	FdPl - Pinegrass - Twinflower		15.0						18.0			21.0			21.0	
04	FdPl - Sitka alder - Pinegrass		18.0			12.0		11	22.6	0.9		21.0		16	20.5	0.9
05	Sxw - Gooseberry - Sarsaparilla		21.0			18.0			21.0			24.0			21.0	
06	Sxw - Oak fern		21.0			18.0			24.0			24.0			24.0	
07	Sxw - Horsetail		18.0			15.0									21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Sx			Py		
		#	SI	SE	#	SI	SE
01	CwSxw - Falsebox	12	19.4	1.2			
02	Fd - Penstemon - Pinegrass					15.0	
03	FdPl - Pinegrass - Twinflower		18.0			18.0	
04	FdPl - Sitka alder - Pinegrass		18.0				
05	Sxw - Gooseberry - Sarsaparilla		21.0				
06	Sxw - Oak fern		21.0				
07	Sxw - Horsetail		18.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		18.0			18.0		7	24.0	0.7		18.0			24.0	
02	Pl - Juniper - Twinflower		15.0			12.0			18.0			12.0			18.0	
03	HwCw - Falsebox - Pipecleaner moss		18.0			15.0			21.0			15.0			21.0	
04	CwFd - Soopolallie - Douglas maple		18.0			15.0			21.0			15.0			21.0	
05	CwHw - Devil's club - Lady fern		21.0			18.0			24.0			18.0			24.0	
06	CwHw - Oval-leaved blueberry - Oak fern		21.0			18.0			24.0			18.0			24.0	
07	CwHw - Horsetail		18.0			15.0						15.0			21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pw			Sx		
		#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		24.0			21.0	
02	Pl - Juniper - Twinflower		18.0			15.0	
03	HwCw - Falsebox - Pipecleaner moss		21.0			18.0	
04	CwFd - Soopolallie - Douglas maple		21.0			18.0	
05	CwHw - Devil's club - Lady fern		24.0			21.0	
06	CwHw - Oval-leaved blueberry - Oak fern		21.0			21.0	
07	CwHw - Horsetail					18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Cw			Fd			Hw			Lw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		18.0			18.0			24.0		9	19.3	1.3		24.0	
03	FdCw - Falsebox - Prince's pine		15.0			15.0		17	22.4	0.8	14	19.4	1.0	16	21.1	0.5
04	CwFd - Falsebox		18.0			18.0			21.0			18.0		10	22.3	1.2
05	CwHw - Oak fern - Foamflower		21.0			18.0			24.0			18.0			24.0	
06	CwHw - Devil's club - Lady fern		21.0			27.0			27.0			21.0			27.0	
07	CwHw - Horsetail		18.0									15.0				
08	CwSxw - Skunk cabbage		18.0									15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Pw			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		24.0			24.0			18.0	
03	FdCw - Falsebox - Prince's pine	11	21.2	0.9		24.0			15.0	
04	CwFd - Falsebox		24.0			24.0			18.0	
05	CwHw - Oak fern - Foamflower		24.0			27.0			24.0	
06	CwHw - Devil's club - Lady fern		27.0			27.0			24.0	
07	CwHw - Horsetail								21.0	
08	CwSxw - Skunk cabbage								18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		18.0			15.0			21.0		9	19.4	0.7		21.0	
01-YC	HwCw - Feathermoss		18.0			15.0			21.0			18.0			21.0	
02	Fd - Juniper - Cladina								15.0						18.0	
03	FdPl - Pinegrass - Feathermoss		15.0			15.0			18.0						21.0	
04	CwFd - Soopolallie - Twinflower					15.0			21.0			15.0			21.0	
05	CwFd - Falsebox		18.0			15.0			21.0			18.0			21.0	
06	CwHw - Oak fern		21.0			18.0			24.0			18.0			24.0	
07	CwHw - Devil's club - Lady fern		21.0			18.0			27.0			18.0				
08	CwSxw - Skunk cabbage		15.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pw			Sx		
		#	SI	SE	#	SI	SE
01	HwCw - Falsebox - Feathermoss		24.0			18.0	
01-YC	HwCw - Feathermoss		24.0			18.0	
02	Fd - Juniper - Cladina						
03	FdPl - Pinegrass - Feathermoss					15.0	
04	CwFd - Soopolallie - Twinflower		21.0			18.0	
05	CwFd - Falsebox		21.0			18.0	
06	CwHw - Oak fern		24.0			21.0	
07	CwHw - Devil's club - Lady fern		27.0			24.0	
08	CwSxw - Skunk cabbage					18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pw		
		#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE
01	CwHw - Devil's club - Lady fern		21.0			21.0			24.0			18.0			24.0	
03	HwCw - Falsebox - Feathermoss		18.0			18.0			24.0			18.0			24.0	
04	CwHw - Oak fern - Spiny wood fern		21.0			18.0			27.0			18.0			27.0	
0	CwSxw - Devil's club - Horsetail		18.0			15.0						15.0			24.0	
0	CwSxw - Skunk cabbage		15.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Sx		
		#	SI	SE
01	CwHw - Devil's club - Lady fern		24.0	
03	HwCw - Falsebox - Feathermoss		21.0	
04	CwHw - Oak fern - Spiny wood fern		24.0	
05	CwSxw - Devil's club - Horsetail		21.0	
06	CwSxw - Skunk cabbage		18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		21.0		10	19.0	1.1		24.0		15	20.3	0.7		24.0	
04	HwCw - Falsebox - Feathermoss		18.0			18.0		8	23.7	1.7		18.0			21.0	
05	CwHw - Devil's club - Lady fern		21.0			21.0			24.0			18.0				
06	CwSxw - Devil's club - Horsetail		18.0			15.0						15.0				
07	Act - Dogwood - Twinberry		18.0			15.0						15.0				
08	CwSxw - Skunk cabbage		18.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pw			Sx		
		#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0	
04	HwCw - Falsebox - Feathermoss		24.0			21.0	
05	CwHw - Devil's club - Lady fern		27.0			24.0	
06	CwSxw - Devil's club - Horsetail					21.0	
07	Act - Dogwood - Twinberry		24.0			21.0	
08	CwSxw - Skunk cabbage					18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bg			Cw			Fd			Hw			Lw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwFd - Mock-orange		18.0			15.0			21.0			18.0			21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Pw			Py		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwFd - Mock-orange		21.0			21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Lw			Pl			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Twinflower	11	18.3	1.0		18.0			18.0			18.0			15.0	
03	FdPy - Bluebunch wheatgrass - Pinegrass		12.0						12.0			15.0				
04	Fd - Pinegrass - Kinnikinnick		15.0			15.0			15.0			18.0				
05	FdLw - Spruce - Pinegrass		21.0			21.0			21.0			21.0			18.0	
06	SxwFd - Dogwood - Gooseberry		24.0			24.0			24.0						24.0	
07	Sxw - Horsetail		21.0			21.0			21.0						21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl		
		#	SI	SE
01	FdPl - Pinegrass - Twinflower			
03	FdPy - Bluebunch wheatgrass - Pinegrass			
04	Fd - Pinegrass - Kinnikinnick			
05	FdLw - Spruce - Pinegrass		18.0	
06	SxwFd - Dogwood - Gooseberry		21.0	
07	Sxw - Horsetail		18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Lw			Pl			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdPl - Pinegrass - Twinflower	14	19.3	1.0		18.0		10	19.9	0.7		18.0				
03	Fd - Snowberry - Balsamroot		15.0			15.0			15.0			15.0				
04	FdLw - Spruce - Pinegrass	8	20.8	0.8		18.0		7	21.7	0.4		18.0			18.0	
05	SxwAt - Sarsaparilla		21.0			21.0			21.0						21.0	
07	Sxw - Horsetail		18.0						18.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Py		
		#	SI	SE	#	SI	SE
01	FdPy - Pinegrass	13	16.3	1.1	8	16.8	1.5
02	FdPy - Bluebunch wheatgrass - Balsamroot		12.0			12.0	
03	FdPy - Bluebunch wheatgrass - Pinegrass		12.0			15.0	
04	FdPy - Snowbrush - Pinegrass		15.0			15.0	
05	FdPy - Pinegrass - Idaho fescue		15.0			15.0	
06	FdPy - Spirea - Feathermoss		18.0			18.0	
07	FdPy - Snowberry - Spirea		18.0			18.0	
08	SxwFd - Douglas maple - Dogwood		21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Fd			Lw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Soopolallie - Grouseberry	9	18.0	0.6	15	20.2	0.6	7	18.8	0.8	10	21.5	0.9	13	19.9	0.5
03	Pl - Juniper - Pinegrass		12.0		10	18.5	0.8		18.0		9	16.8	1.3		15.0	
04	Pl - Oregon-grape - Pinegrass	7	18.4	1.3	31	19.0	0.5	10	16.4	0.6	18	18.1	0.7	7	22.7	1.9
05	Sxw - Soopolallie - Snowberry		15.0		7	17.3	0.6	7	21.2	1.2	7	19.6	0.6	7	22.4	1.2
06	Sxw - Dogwood - Horsetail		15.0			18.0						15.0			18.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	At		
		#	SI	SE
01	Sxw - Soopolallie - Grouseberry			
03	Pl - Juniper - Pinegrass	7	19.5	1.6
04	Pl - Oregon-grape - Pinegrass			
05	Sxw - Soopolallie - Snowberry			
06	Sxw - Dogwood - Horsetail			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] MS...

Site Series(*)	Name	Bl			Fd			Lw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Falsebox - Feathermoss	10	19.3	1.0		18.0			21.0		22	19.2	0.6	14	17.5	1.2
02	Fd - Penstemon - Pinegrass					15.0			15.0			15.0				
03	Pl - Grouseberry - Cladonia		15.0			18.0			18.0		7	13.3	1.7		15.0	
04	Pl - Pinegrass - Kinnikinnick		15.0			18.0			21.0		11	17.6	1.2		15.0	
05	Sxw - Trapper's tea - Grouseberry		18.0			18.0					8	19.7	0.6	7	16.3	0.4
06	Sxw - Gooseberry		18.0			18.0						21.0			18.0	
07	Sxw - Trapper's tea - Horsetail		15.0									18.0			15.0	
08	Sxw - Gooseberry - Oak fern		18.0			18.0						21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Fd			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Py - Bluebunch wheatgrass - Junegrass		15.0			15.0				
04	FdPy - Ninebark		18.0			18.0				
05	PyAct - Snowberry - Kentucky bluegrass		18.0			18.0			18.0	
06	AtAct - Snowberry - Horsetail		21.0						21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd			Lw			Pl			Py			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Py - Bluebunch wheatgrass - Junegrass		15.0			12.0			12.0			15.0				
03	PyAt - Rose - Solomon's-seal		18.0			15.0			15.0			18.0			15.0	
04	Act - Dogwood - Nootka rose														18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

MBSdk1

Site Series(*)	Name	Pl			Sw			Sb			Bl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Knight's plume - Step moss	15	15.4	0.5	8	14.0	0.8						
02	Pl - Lingonberry - Feathermoss		12.0			9.0							
03	Sw - Wildrye - Toad-flax		12.0			12.0							
04	Sb - Lingonberry - Knight's plume	15	13.7	0.4		12.0			9.0				
05	SwPl - Soopolallie - Twinflower		18.0			15.0						12.0	
06	Sw - Scouring-rush - Step moss		15.0			12.0						12.0	
07	Sb - Lingonberry - Coltsfoot		12.0			12.0			12.0				
08	Sw - Currant - Horsetail		18.0			18.0							
09	Sb - Horsetail - Sphagnum		9.0			9.0							
10	Sb - Labrador tea - Sphagnum		6.0			6.0							
11	Sw - Willow - Glow moss		6.0			6.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] WBSdk2

Site Series(*)	Name	Pl			Sb			Sw			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Knight's plume - Step moss	15	12.8	0.6		9.0			12.0				
02	Pl - Lingonberry - Cladonia	8	11.4	0.6					9.0				
03	Sb - Lingonberry - Knight's plume		9.0			6.0			9.0				
04	Sb - Labrador tea - Feathermoss		9.0			6.0			9.0				
05	Sw - Wildrye - Toad-flax	12	14.7	0.8					12.0				
06	Sw - Currant - Horsetail		15.0						15.0				
07	Sb - Labrador tea - Sphagnum		6.0						6.0				
08	Lt - Glow moss								9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] WBSmw1

Site Series(*)	Name	At			Pl			Sw			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SwAt - Step moss	13	17.6	0.8		18.0		16	17.7	0.7			
02	Pl - Lingonberry - Velvet-leaved blueberry					12.0			9.0				
03	Sw - Wildrye - Peavine	13	16.5	0.7		18.0		9	17.0	0.6			
04	Sb - Lingonberry - Coltsfoot					15.0			12.0			9.0	
05	Sw - Currant - Oak fern					18.0			18.0			15.0	
06	Sw - Currant - Bluebells	8	17.1	1.2		18.0		9	18.1	0.9			
07	Sw - Currant - Horsetail					18.0			18.0				
08	Sb - Labrador tea - Sphagnum					12.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

MBSmw2

Site Series(*)	Name	Pl			Sw			Sb			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SwAt - Step moss		15.0			15.0							
02	Pl - Lingonberry - Velvet-leaved blueberry		12.0			9.0							
03	Sb - Lingonberry - Knight's plume		15.0			12.0			12.0				
04	Sb - Lingonberry - Coltsfoot		15.0			12.0			12.0				
05	Sw - Currant - Horsetail					18.0							
06	Sb - Feathermoss - Bluebells		18.0			15.0		7	8.4	0.9			
07	Lt - Horsetail					12.0			15.0			15.0	
08	Sb - Cloudberry - Sphagnum		12.0			9.0		8	8.9	0.5			
09	Sb - Willow - Glow moss					9.0		8	12.4	0.9		12.0	
10	Lt - Buckbean		12.0			9.0			9.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

MBSwk1

Site Series(*)	Name	Pl			Sw			Sb			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Huckleberry - Step moss		15.0			12.0							
02	Pl - Lingonberry - Velvet-leaved blueberry		12.0			9.0							
03	Sb - Lingonberry - Coltsfoot		12.0			9.0			9.0				
04	Sw - Wildrye - Peavine		15.0			12.0							
05	Sw - Currant - Bluebells		15.0			15.0							
06	Sw - Currant - Horsetail		15.0			15.0							
07	Sb - Horsetail - Sphagnum		12.0			9.0			9.0				
08	Sb - Willow - Glow moss		9.0			6.0			6.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

WBSwk2

Site Series(*)	Name	Pl			Sw			Sb			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Huckleberry - Step moss		15.0			12.0							
02	Pl - Lingonberry - Feathermoss		12.0			9.0							
03	Sw - Wildrye - Peavine		15.0			12.0							
04	Sb - Lingonberry - Coltsfoot		12.0			9.0			9.0				
05	Sw - Currant - Bluebells		15.0			15.0							
06	Sw - Currant - Horsetail		15.0			15.0							
07	Sb - Horsetail - Sphagnum		12.0			9.0			9.0				
08	Sb - Willow - Glow moss		9.0			6.0			6.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] MBSwk3

Site Series(*)	Name	Pl			Sw		
		#	SI	SE	#	SI	SE
01	Sw - Huckleberry - Step moss		12.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Gooseberry		15.0			18.0			15.0	
02	Bl - Huckleberry - Feathermoss		6.0			9.0			9.0	
03	BlPl - Cladina		12.0			12.0			12.0	
04	Bl - Azalea - Rhododendron		12.0			15.0			12.0	
05	Bl - Oak fern- Bramble		15.0			15.0			15.0	
06	Bl - Devil's club - Lady fern		15.0			18.0			18.0	
07	Bl - Labrador tea - Horsetail		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SFmv1

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Feathermoss		12.0			15.0			15.0	
02	Pl - Huckleberry - Cladonia		9.0			12.0				
03	Bl - Huckleberry - Feathermoss		9.0			12.0			12.0	
04	Bl - Huckleberry - Gooseberry		15.0			18.0			15.0	
05	Bl - Horsetail - Glow moss		15.0			15.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]SFmv2

Site Series(*)	Name	Bl			Pl			Se			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Feathermoss		12.0			15.0			15.0				
02	Bl - Lingonberry		9.0			12.0			9.0				
03	BlSb - Labrador tea		6.0			9.0			6.0			6.0	
04	Bl - Oak fern - Knight's plume		15.0			18.0			15.0				
05	Bl - Devil's club - Rhododendron		15.0			15.0			15.0				
06	Bl - Alder - Horsetail		15.0			15.0			15.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

ESSFmv3

Site Series(*)	Name	Bl			Pl			Se			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Feathermoss		12.0			12.0			12.0				
02	BlPl - Crowberry - Cladina		9.0			9.0			9.0				
03	BlSb - Labrador tea		6.0			9.0			6.0			6.0	
04	Bl - Oak fern - Knight's plume		12.0			15.0			15.0				
05	Bl - Devil's club - Rhododendron		12.0			15.0			15.0				
06	Sxw - Huckleberry - Highbush-cranberry		12.0			12.0							
07	Bl - Horsetail - Feathermoss		9.0			9.0			9.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SSFmv4

Site Series(*)	Name	Bl			Pl			Se			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Feathermoss		12.0			15.0			15.0				
02	BlPl - Crowberry - Cladina		9.0			12.0			9.0				
03	BlSb - Labrador tea		6.0			9.0			6.0			6.0	
04	Bl - Rhododendron - Horsetail		15.0			18.0			15.0				
05	Bl - Alder - Horsetail		15.0			15.0			15.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SSFwc2

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Azalea - Oak fern		15.0			15.0			15.0	
02	Pl - Huckleberry - Cladonia		9.0			12.0			12.0	
03	Bl - Huckleberry - Arnica		12.0			15.0			12.0	
04	Bl - Rhododendron - Heron's-bill		12.0			15.0			12.0	
05	Bl - Azalea - Feathermoss	11	13.6	0.8		15.0			12.0	
06	Bl - Valerian - Oak fern		15.0			18.0			15.0	
07	Bl - Devil's club - Lady fern		18.0			18.0			21.0	
08	Bl - Horsetail - Sphagnum		15.0			12.0			15.0	
09	Pl - Dwarf blueberry - Sphagnum		12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SSFwc3

Site Series(*)	Name	Bl			Se		
		#	Sl	SE	#	Sl	SE
01	Bl - Rhododendron - Oak fern		15.0			15.0	
02	Bl - Rhododendron - Queen's cup		9.0			9.0	
03	Bl - Globeflower - Horsetail		15.0			15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SSFwk1

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Oak fern - Brachythecium		15.0			18.0			15.0	
02	Bl - Huckleberry - Feathermoss		9.0			12.0			9.0	
03	Bl - Oak fern - Knight's plume		12.0			15.0			12.0	
04	Bl - Twinberry - Lady fern		15.0			18.0			15.0	
05	Bl - Devil's club - Lady fern		15.0			18.0			15.0	
06	Bl - Horsetail - Sphagnum		9.0						9.0	
07	Bl - Lady fern - Horsetail		12.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]SFwk2

Site Series(*)	Name	Bl			Se			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Oak fern - Knight's plume		15.0			15.0				
02	Bl - Oak fern - Sarsaparilla		9.0			9.0				
03	Bl - Oak fern - Bluebells		12.0			12.0			15.0	
04	Bl - Devil's club - Rhododendron		15.0			15.0				
05	Bl - Rhododendron - Lady fern		15.0			15.0				
06	Bl - Horsetail - Sphagnum		12.0			12.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Spruce - Step moss	8	20.5	1.4	7	17.3	1.4	15	22.9	0.9	7	20.8	2.5	7	23.6	1.0
02	CwSxw - Soopolallie		12.0			12.0			15.0			12.0			15.0	
03	HwCw - Step moss	7	20.3	1.5		12.0		8	21.9	1.2		15.0		10	22.0	0.7
04	CwHw - Oak fern		15.0			15.0			21.0			18.0			21.0	
05	CwHw - Devil's club - Lady fern		18.0			18.0			24.0			18.0			24.0	
06	CwSxw - Devil's club - Horsetail		18.0			18.0						18.0			21.0	
07	SbPl - Bog-laurel - Sphagnum														15.0	
08	Skunk cabbage - Sphagnum		18.0			18.0									15.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Sx			Pw			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwCw - Spruce - Step moss	13	22.8	0.7						
02	CwSxw - Soopolallie		12.0							
03	HwCw - Step moss		15.0							
04	CwHw - Oak fern		18.0			21.0				
05	CwHw - Devil's club - Lady fern		21.0							
06	CwSxw - Devil's club - Horsetail		21.0							
07	SbPl - Bog-laurel - Sphagnum		12.0						12.0	
08	Skunk cabbage - Sphagnum		18.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] ttvk2

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Devil's club - Lady fern		15.0			18.0			21.0			15.0			24.0	
02	HwCw - Cladonia					12.0						12.0			15.0	
03	HwCw - Step moss		12.0			12.0			15.0			12.0			15.0	
04	CwHw - Oak fern		15.0			12.0			18.0			15.0			18.0	
05	Cw - Devil's club - Ostrich fern		15.0			18.0						18.0			21.0	
06	CwSxw - Skunk cabbage		18.0			18.0						15.0			18.0	
07	Sb - Sphagnum														15.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted].k2

Site Series(*)	Name	Sx			Sb		
		#	SI	SE	#	SI	SE
01	CwHw - Devil's club - Lady fern		21.0				
02	HwCw - Cladonia		12.0				
03	HwCw - Step moss		12.0				
04	CwHw - Oak fern		15.0				
05	Cw - Devil's club - Ostrich fern		21.0				
06	CwSxw - Skunk cabbage		18.0				
07	Sb - Sphagnum		12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

wk1

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		21.0		10	19.0	1.1		24.0		15	20.3	0.7		24.0	
03	PlHw - Velvet-leaved blueberry		18.0			15.0			21.0			15.0			21.0	
04	HwCw - Falsebox - Feathermoss		18.0			18.0		8	23.7	1.7		18.0			21.0	
05	CwHw - Devil's club - Lady fern		21.0			21.0			24.0			18.0				
06	CwSxw - Devil's club - Horsetail		18.0			15.0						15.0				
07	Act - Dogwood - Twinberry					15.0						15.0				
08	CwSxw - Skunk cabbage		18.0			15.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted].wk1

Site Series(*)	Name	Pw			Sx		
		#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0	
03	PlHw - Velvet-leaved blueberry		24.0			18.0	
04	HwCw - Falsebox - Feathermoss		24.0			21.0	
05	CwHw - Devil's club - Lady fern		27.0			24.0	
06	CwSxw - Devil's club - Horsetail					21.0	
07	Act - Dogwood - Twinberry		24.0			21.0	
08	CwSxw - Skunk cabbage					18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] wk3

Site Series(*)	Name	Bl			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		15.0			18.0		10	23.5	0.7	9	19.7	1.1		21.0	
02	Hw - Azalea - Cladonia											12.0			15.0	
03	CwSxw - Prince's pine - Cat's-tail moss		12.0			12.0			15.0			12.0			15.0	
04	HwCw - Step moss		15.0			15.0		7	21.9	1.6		15.0			18.0	
05	CwHw - Devil's club - Lady fern		18.0			18.0			21.0			18.0			24.0	
06	CwSxw - Devil's club - Horsetail		18.0			18.0						18.0			24.0	
07	Hw - Wood horsetail - Sphagnum											15.0			18.0	
08	PlSb - Sedge - Sphagnum														15.0	
09	CwSxw - Skunk cabbage		18.0			18.0						15.0				

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] wk3

Site Series(*)	Name	Sx			Sb		
		#	SI	SE	#	SI	SE
01	CwHw - Oak fern	24	22.0	0.6			
02	Hw - Azalea - Cladonia		12.0				
03	CwSxw - Prince's pine - Cat's-tail moss		12.0				
04	HwCw - Step moss		15.0				
05	CwHw - Devil's club - Lady fern	8	19.8	1.1			
06	CwSxw - Devil's club - Horsetail		21.0				
07	Hw - Wood horsetail - Sphagnum		15.0				
08	PlSb - Sedge - Sphagnum					12.0	
09	CwSxw - Skunk cabbage		18.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] wk4

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Oak fern		24.0			24.0			21.0	
02	HwCw - Cladonia		15.0			15.0				
03	CwSxw - Soopolallie		15.0			15.0				
04	CwSxw - Velvet-leaved blueberry		18.0			21.0			18.0	
05	HwCw - Step moss		21.0			21.0			18.0	
06	Sxw - Twinberry - Oak fern					21.0			21.0	
07	CwHw - Devil's club - Lady fern		24.0			21.0			24.0	
08	Sxw - Devil's club - Lady fern								18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]SBPSmc

Site Series(*)	Name	Pl			Sb			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Feathermoss - Cladina		12.0			9.0			9.0	
02	Pl - Kinnikinnick - Cladonia		9.0						6.0	
03	SbPl - Feathermoss		9.0						6.0	
04	Sxw - Scrub birch - Feathermoss		9.0						6.0	
05	Sxw - Horsetail		15.0						12.0	
06	Sxw - Horsetail - Glow moss		12.0						9.0	
07	SbSxw - Scrub birch - Sedge		9.0						9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]BSdh

Site Series(*)	Name	Fd			Pl			Sx			Lw			Pw		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Ricegrass		21.0			21.0			21.0							
02	Pl - Velvet-leaved blueberry - Cladonia					15.0			12.0							
03	FdPl - Pinegrass - Feathermoss		15.0			18.0						18.0			18.0	
04	Pl - Pinegrass - Feathermoss		15.0			18.0			15.0							
05	Pl - Labrador tea - Velvet-leaved blueberry					18.0			15.0							
06	SxwFd - Thimbleberry					21.0			21.0							
07	Sxw - Horsetail					21.0			21.0							
08	Sb - Scrub birch - Sedge					15.0			12.0							

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SBSdh

Site Series(*)	Name	Sb		
		#	SI	SE
01	SxwFd - Ricegrass			
02	Pl - Velvet-leaved blueberry - Cladonia			
03	FdPl - Pinegrass - Feathermoss			
04	Pl - Pinegrass - Feathermoss			
05	Pl - Labrador tea - Velvet-leaved blueberry		15.0	
06	SxwFd - Thimbleberry			
07	Sxw - Horsetail			
08	Sb - Scrub birch - Sedge		12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

BSdh1

Site Series(*)	Name	Fd		
		#	SI	SE
01	SxwFd - Ricegrass	9	21.9	1.5

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSdk

Site Series(*)	Name	Pl			Sx			Sb			Fd		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Spirea - Purple peavine	24	19.5	0.4	17	18.0	0.6						
02	Pl - Juniper - Ricegrass		15.0			15.0							
03	Pl - Feathermoss - Cladina	13	16.2	0.6	10	14.9	0.6		9.0				
04	Fd - Soopolallie - Feathermoss		15.0			15.0							
05	Sxw - Spirea - Feathermoss	9	18.9	1.0		18.0							
06	Sxw - Twinberry - Coltsfoot	14	21.9	0.6	22	20.4	0.6					18.0	
07	Sxw - Horsetail		21.0			21.0							
08	Act - Dogwood - Prickly rose		21.0			21.0							
09	Sb - Creeping-snowberry - Sphagnum		12.0						9.0				
10	Sb - Soft-leaved sedge - Sphagnum		12.0			12.0			9.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]BSdw1

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass		21.0			24.0			21.0	
02	FdPl - Cladonia		15.0			15.0				
03	Fd - Saskatoon - Pinegrass		15.0			15.0				
04	Pl - Pinegrass - Feathermoss		15.0			18.0			15.0	
05	SxwFd - Ricegrass		21.0			21.0			18.0	
06	SxwFd - Thimbleberry		21.0			24.0			21.0	
07	Sxw - Twinberry - Coltsfoot		21.0			21.0			21.0	
08	Sxw - Twinberry - Oak fern		21.0			24.0			24.0	
09	Sxw - Horsetail - Glow moss					21.0			21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Subzone: SBSdw2

Site Series(*)	Name	Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass		18.0			21.0			18.0	
02	FdPl - Cladonia		15.0			15.0				
03	Pl - Kinnikinnick - Wavy-leaved moss					15.0				
04	Fd - Pinegrass - Aster		15.0			15.0				
05	SxwFd - Cat's-tail moss		18.0			21.0			18.0	
06	Pl - Pinegrass - Feathermoss		18.0			21.0			18.0	
07	PlSb - Feathermoss					18.0			15.0	
08	Sxw - Twinberry		18.0			21.0			18.0	
09	Sxw - Devil's club - Knight's plume		21.0			21.0			21.0	
10	Sxw - Horsetail					15.0			18.0	
11	Sb - Soft-leaved sedge - Sphagnum					9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[illegible] SBSdw3

Site Series(*)	Name	At			Ep			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass	9	20.1	1.2	7	18.6	1.2		18.0		13	21.5	0.7	15	18.4	0.8
02	FdPl - Cladonia								15.0			15.0				
03	Pl - Feathermoss - Cladina											15.0			12.0	
04	SxwFd - Ricegrass							10	18.5	0.7		18.0			15.0	
05	PlSb - Feathermoss											15.0			12.0	
06	Sxw - Pink spirea - Prickly rose											18.0			15.0	
07	Sxw - Twinberry											21.0			18.0	
08	Sxw - Oak fern								18.0			21.0			18.0	
09	Sxw - Horsetail - Glow moss											18.0			18.0	
10	Sb - Soft-leaved sedge - Sphagnum											15.0			12.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSdw3

Site Series(*)	Name	Sb			Bl			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Pinegrass									
02	FdPl - Cladonia									
03	Pl - Feathermoss - Cladina									
04	SxwFd - Ricegrass									
05	PlSb - Feathermoss		12.0							
06	Sxw - Pink spirea - Prickly rose					15.0				
07	Sxw - Twinberry					15.0				
08	Sxw - Oak fern					15.0				
09	Sxw - Horsetail - Glow moss									
10	Sb - Soft-leaved sedge - Sphagnum		12.0						15.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SBSmc2

Site Series(*)	Name	Bl			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry	31	15.4	0.5	97	17.9	0.3	48	18.7	0.5			
02	Pl - Huckleberry - Cladonia		12.0		27	16.4	0.5	7	12.3	0.8			
03	SbPl - Feathermoss					15.0			15.0			9.0	
04	Sxw - Huckleberry - Dwarf blueberry					18.0			18.0				
05	Sxw - Twinberry - Coltsfoot	11	16.3	1.0	26	19.1	0.5	35	21.3	0.6			
06	Sxw - Oak fern	17	19.1	0.7	10	19.9	0.5	26	19.3	0.5			
07	Sxw - Scrub birch - Feathermoss					12.0			12.0			9.0	
08	Sxw - Twinberry - Oak fern					21.0			21.0				
09	Sxw - Devil's club		18.0			21.0		8	19.8	0.8			
10	Sxw - Horsetail	8	15.3	0.9	7	19.6	0.8	9	19.3	1.0			
11	Sxw - Horsetail - Glow moss					18.0			18.0				
12	SbSxw - Scrub birch - Sedge					12.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSmc3

Site Series(*)	Name	Bl			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry		15.0			18.0			15.0				
02	Pl - Juniper - Dwarf blueberry					15.0			12.0				
03	Pl - Feathermoss - Cladina					15.0			12.0				
04	Sxw - Huckleberry - Soopolallie		12.0			15.0			12.0				
05	Sb - Huckleberry - Spirea					15.0			12.0			12.0	
06	SbPl - Feathermoss					15.0			12.0			12.0	
07	Sxw - Twinberry		15.0			18.0			15.0				
08	Sxw - Horsetail		15.0			18.0			15.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSmh

Site Series(*)	Name	Fd			Sx			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Hazelnut		21.0			18.0				
02	FdPl - Cladonia		18.0							
03	FdPl - Velvet-leaved blueberry - Cladonia		18.0			15.0			21.0	
04	Fd - Douglas maple - Step moss		21.0			15.0				
05	SxwFd - Feathermoss		21.0			18.0				
06	SxwFd - Coltsfoot		21.0			21.0				
07	SxwEp - Devil's club		24.0			24.0				
08	Sxw - Ostrich fern		24.0			24.0				
09	Sxw - Horsetail - Glow moss					21.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSmk1

Site Series(*)	Name	At			Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry - Highbush-cranberry	11	19.4	1.1	12	15.7	1.2		21.0		19	20.1	0.7	20	19.1	0.8
02	Pl - Cladina - Step moss											18.0			15.0	
03	Pl - Feathermoss - Cladina										10	16.1	0.9		15.0	
04	SxwFd - Knight's plume					15.0			18.0			21.0			18.0	
05	SxwFd - Toad-flax					18.0		7	21.2	1.2	14	20.9	1.2		18.0	
06	Sb - Huckleberry - Spirea										7	17.6	1.0		15.0	
07	Sxw - Oak fern					18.0			21.0			21.0		9	20.8	1.5
08	Sxw - Devil's club					18.0						21.0			18.0	
09	Sxw - Horsetail					15.0						21.0			18.0	
10	Sb - Scrub birch - Sedge											15.0			12.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

SBSmk1

Site Series(*)	Name	Sb		
		#	SI	SE
01	Sxw - Huckleberry - Highbush-cranberry			
02	Pl - Cladina - Step moss			
03	Pl - Feathermoss - Cladina			
04	SxwFd - Knight's plume			
05	SxwFd - Toad-flax			
06	Sb - Huckleberry - Spirea		15.0	
07	Sxw - Oak fern			
08	Sxw - Devil's club			
09	Sxw - Horsetail			
10	Sb - Scrub birch - Sedge		12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]BSmk2

Site Series(*)	Name	Bl			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry - Highbush-cranberry		15.0			21.0			18.0				
02	Pl - Feathermoss - Cladina					18.0			15.0				
03	Sxw - Huckleberry - Soopolallie		12.0			18.0			15.0				
04	Sb - Huckleberry - Spirea					18.0			12.0			15.0	
05	Sxw - Oak fern		18.0			21.0			21.0				
06	Sxw - Horsetail		18.0			21.0			21.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]BSmw

Site Series(*)	Name	Bl			Fd			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	SxwFd - Falsebox		15.0		8	22.6	0.6	15	22.4	0.6	22	20 0	0 4
02	FdBl - Huckleberry					15.0			15.0			15 0	
03	Pl - Huckleberry - Velvet-leaved blueberry					18.0		8	17.1	1.2		9.0	
04	SxwFd - Knight's plume					18.0			21.0			15.0	
05	Sxw - Pink spirea							7	18.2	0.7		15.0	
06	Sxw - Oak fern		15.0			21.0			24.0			21.0	
07	Sxw - Twinberry - Oak fern		15.0						24.0			21.0	
08	Sxw - Devil's club		18.0			21.0			24.0			24.0	
09	Sxw - Horsetail		15.0						18.0			18.0	
10	SbSxw - Scrub birch - Sedge								9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Fd			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Devil's club		15.0			18.0			21.0		8	19.7	1.3			
02	Pl - Huckleberry - Velvet-leaved blueberry		12.0						18.0			15.0				
03	SxwFd - Thimbleberry		12.0			15.0			18.0			15.0				
04	Sxw - Oak fern		15.0			18.0			21.0			18.0				
05	Sxw - Devil's club - Spiny wood fern		18.0						21.0			18.0				
06	Sxw - Horsetail		18.0						18.0			18.0				
07	Sxw - Devil's club - Ostrich fern		15.0						18.0			18.0				
08	SbPl - Bog-laurel - Sphagnum								12.0			12.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

wk1

Site Series(*)	Name	Bl			Fd			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Oak fern		15.0			21.0			21.0		13	21.2	0.8			
02	Pl - Huckleberry - Cladina		12.0			15.0			15.0			12.0				
03	Pl - Huckleberry - Velvet-leaved blueberry		12.0			15.0			18.0			15.0				
04	SxwFd - Knight's plume		15.0			18.0			18.0			15.0				
05	Sxw - Huckleberry - Highbush-cranberry		15.0			18.0			21.0			18.0				
06	Sxw - Pink spirea - Oak fern		15.0						21.0			18.0				
07	Sxw - Twinberry - Oak fern		18.0						21.0			18.0				
08	Sxw - Devil's club		18.0						21.0			18.0				
09	Sxw - Horsetail		18.0						21.0			18.0				
10	Sxw - Devil's club - Lady fern		18.0						21.0			18.0				
11	SbSxw - Scrub birch - Sedge								12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] wk2

Site Series(*)	Name	Bl			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Oak fern		15.0			21.0		12	21.8	0.9			
02	Pl - Huckleberry - Cladina		12.0			15.0			15.0				
03	Sxw - Huckleberry - Highbush-cranberry		12.0			18.0			18.0				
04	SbPl - Feathermoss					18.0			15.0			15.0	
05	Sxw - Devil's club		18.0			21.0			21.0				
06	Sxw - Horsetail		18.0			21.0			24.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]wk3

Site Series(*)	Name	Bl			Pl			Sx			Fd			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Oak fern	7	15.3	1.1	12	20.6	0.9	13	16.7	0.6						
02	Pl - Huckleberry - Cladina		12.0			15.0			12.0							
03	SxwFd - Purple peavine					15.0			15.0			15.0				
04	Sxw - Huckleberry - Highbush-cranberry		15.0			18.0			15.0							
05	Sb - Labrador tea				7	15.7	1.1		15.0					7	11.2	1.2
06	Sxw - Twinberry - Coltsfoot		15.0		7	20.7	0.8	7	19.2	1.4						
07	Sxw - Devil's club		18.0			21.0			21.0							
08	Sxw - Horsetail		18.0			21.0			18.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] WBSdk1

Site Series(*)	Name	Bl			Pl			Sw			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Knight's plume - Step moss		12.0		15	15.4	0.5	8	14.0	0.8			
02	Pl - Lingonberry - Feathermoss					12.0			9.0				
03	Sw - Wildrye - Toad-flax					12.0			12.0				
04	Sb - Lingonberry - Knight's plume				15	13.7	0.4		12.0			9.0	
05	Sw - Soopolallie - Twinflower		12.0			18.0			15.0				
06	Sw - Scouring-rush - Step moss		12.0			15.0			15.0				
07	Sb - Lingonberry - Coltsfoot					12.0			12.0			12.0	
08	Sw - Currant - Horsetail					18.0			18.0				
09	Sb - Horsetail - Sphagnum					9.0			9.0				
10	Sb - Labrador tea - Sphagnum					6.0			6.0				
11	Sw - Willow - Glow moss					6.0			6.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]WBSdk2

Site Series(*)	Name	Pl			Sb			Sw			Lt		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sw - Knight's plume - Step moss	15	12.8	0.6		9.0			12.0				
02	Pl - Lingonberry - Cladonia	8	11.4	0.6					9.0				
03	Sb - Lingonberry - Knight's plume		9.0			6.0			9.0				
04	Sb - Labrador tea - Feathermoss		9.0			6.0			9.0				
05	Sw - Wildrye - Toad-flax	12	14.7	0.8					12.0				
06	Sw - Currant - Horsetail		15.0						15.0				
07	Sb - Labrador tea - Sphagnum		6.0						6.0				
08	Lt - Glow moss								9.0			9 0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] .vrivh2

Site Series(*)	Name	Ba			Cw			Hw			Pl			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Salal		16.0		7	19.7	0.8		20.0			20.0			16.0	
03	CwYc - Salal					16.0			16.0			16.0				
04	HwSs - Lanky moss	33	27.3	0.7	20	22.8	0.5	43	27.0	0.6				43	31.6	0.6
05	CwSs - Sword fern		24.0			24.0			24.0						28.0	
06	CwSs - Foamflower	15	31.0	0.7	10	24.2	0.5	18	28.2	0.7				27	32.9	0.8
07	CwSs - Devil's club		28.0		9	22.0	1.2	9	24.7	2.3				18	30.5	1.5
08	Ss - Lily-of-the-valley		28.0			24.0			24.0						32.0	
09	Ss - Trisetum								28.0						32.0	
11	CwYc - Goldthread					12.0			16.0			16.0				
12	PlYc - Sphagnum					8.0						12.0				
13	CwSs - Skunk cabbage					16.0			16.0						20.0	
14	Ss - Salal					12.0			12.0						16.0	
15	Ss - Kindbergia					16.0			16.0						24.0	
16	Ss - Reedgrass					12.0			12.0						16.0	
17	Ss - Sword fern					24.0			24.0						32.0	
18	Ss - Slough sedge					16.0									20.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] MHvh2

Site Series(*)	Name	Yc		
		#	SI	SE
01	CwHw - Salal		16.0	
03	CwYc - Salal			
04	HwSs - Lanky moss			
05	CwSs - Sword fern			
06	CwSs - Foamflower			
07	CwSs - Devil's club			
08	Ss - Lily-of-the-valley			
09	Ss - Trisetum			
11	CwYc - Goldthread		12.0	
12	PlYc - Sphagnum		8.0	
13	CwSs - Skunk cabbage		16.0	
14	Ss - Salal			
15	Ss - Kindbergia			
16	Ss - Reedgrass			
17	Ss - Sword fern			
18	Ss - Slough sedge			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Cw			Hw			Ss			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry	43	29.3	0.6	34	22.6	0.5	94	27.7	0.4	37	30.8	0.7			
02	HwPl - Cladina					8.0			8.0						12.0	
03	HwCw - Salal					16.0			16.0						16.0	
04	CwHw - Sword fern		24.0		10	22.5	1.2	7	26.2	2.3		24.0				
05	BaCw - Foamflower	27	30.9	0.5	15	24.0	0.8	35	28.5	0.6	30	32.7	0.9			
06	HwBa - Deer fern		24.0		9	23.3	1.9	20	25.2	1.4		28.0				
08	BaSs - Devil's club		28.0		8	22.1	1.2	16	26.3	1.0	29	31.4	0.9			
09	Ss - Salmonberry		28.0			24.0			28.0			32.0				
10	Act - Red-osier dogwood		28.0			24.0						32.0				
12	CwYc - Goldthread					12.0			16.0						16.0	
13	Pl - Sphagnum					8.0									12.0	
14	CwSs - Skunk cabbage				13	19.4	0.9	10	21.0	1.6	8	26.0	1.7			

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Hm			Yc		
		#	SI	SE	#	SI	SE
01	HwBa - Blueberry						
02	HwPl - Cladina						
03	HwCw - Salal						
04	CwHw - Sword fern						
05	BaCw - Foamflower						
06	HwBa - Deer fern						
08	BaSs - Devil's club						
09	Ss - Salmonberry						
10	Act - Red-osier dogwood						
12	CwYc - Goldthread		16.0			12.0	
13	Pl - Sphagnum					8.0	
14	CwSs - Skunk cabbage						

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Cw			Hw			Ss			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry		24.0			20.0			24.0			20.0			20.0	
02	HwPl - Cladina					8.0			8.0						8.0	
03	HwCw - Salal		16.0			16.0			16.0						16.0	
04	CwHw - Sword fern		20.0			20.0			24.0			24.0			20.0	
05	BaCw - Foamflower		28.0			24.0			28.0			24.0			24.0	
06	HwBa - Deer fern		24.0			20.0			24.0			24.0			20.0	
08	BaSs - Devil's club		28.0			24.0			28.0			28.0			24.0	
09	CwYc - Goldthread					12.0			12.0						12.0	
10	Pl - Sphagnum														8.0	
11	CwYc - Skunk cabbage					16.0			16.0			20.0			16.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

WHvm2

Site Series(*)	Name	Pl			Fd			Hm		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry									
02	HwPl - Cladina		12.0							
03	HwCw - Salal		20.0			24.0				
04	CwHw - Sword fern									
05	BaCw - Foamflower									
06	HwBa - Deer fern									
08	BaSs - Devil's club									
09	CwYc - Goldthread		12.0						12.0	
10	Pl - Sphagnum		12.0							
11	CwYc - Skunk cabbage								16.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Wfhwm

Site Series(*)	Name	Ba			Cw			Hm			Hw			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwSs - Blueberry		24.0		9	19.7	0.9		24.0		17	19.6	0.7	10	21.4	1.3
02	HwSs - Step moss					16.0			20.0			20.0			16.0	
03	SsHw - Oak fern		28.0			24.0						28.0		7	30.3	1.7
04	SsHw - Devil's club		28.0			24.0						28.0			32.0	
05	Ss - Salmonberry		28.0			24.0						28.0			32.0	
06	Act - Red-osier dogwood		28.0			24.0									32.0	
08	Hw - Sphagnum					12.0			16.0			16.0				
09	Ss - Skunk cabbage					16.0						16.0			20.0	
10	Pl - Sphagnum					8.0										

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] Mhwm

Site Series(*)	Name	Yc			Pl		
		#	SI	SE	#	SI	SE
01	HwSs - Blueberry		20.0				
02	HwSs - Step moss		16.0			20.0	
03	SsHw - Oak fern						
04	SsHw - Devil's club						
05	Ss - Salmonberry						
06	Act - Red-osier dogwood						
08	Hw - Sphagnum		12.0			16.0	
09	Ss - Skunk cabbage						
10	Pl - Sphagnum		8.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Cw			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Bramble		20.0			16.0		8	22.6	1.3		20.0			20.0	
02	Pl - Kinnikinnick					8.0			12.0			12.0				
03	HwPl - Feathermoss					16.0			16.0			20.0				
04	BaCw - Oak fern		24.0			20.0			24.0			24.0			24.0	
05	HwBa - Queen's cup		20.0			16.0			20.0						24.0	
06	BaCw - Devil's club		28.0			24.0			24.0						28.0	
07	Ss - Salmonberry		28.0			20.0									32.0	
08	Act - Red-osier dogwood		28.0			20.0									32.0	
10	Pl - Sphagnum					8.0						12.0				
11	CwSs - Skunk cabbage		12.0			16.0			12.0						16.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] ws2

Site Series(*)	Name	Ba			Cw			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Bramble		20.0			16.0			20.0			20.0			20.0	
02	Pl - Kinnikinnick					8.0			12.0			12.0				
03	HwPl - Feathermoss					16.0			16.0			16.0				
04	BaCw - Oak fern		24.0			16.0			24.0						24.0	
05	HwBa - Queen's cup		20.0			16.0			20.0						20.0	
06	BaCw - Devil's club		24.0			20.0			24.0						24.0	
07	Ss - Salmonberry		28.0			20.0			28.0						28.0	
08	Act - Red-osier dogwood		28.0			20.0									28.0	
10	Pl - Sphagnum					8.0						12.0				
11	CwSs - Skunk cabbage		12.0			12.0			12.0						12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Huckleberry - Leafy liverwort	10	13.8	1.3	11	16.2	0.7	10	15.4	1.2
02	BlPl - Juniper - Cladonia		9.0			9.0			9.0	
03	Bl - Huckleberry - Crowberry		9.0			9.0			9.0	
04	Bl - Huckleberry - Heron's-bill	10	12.8	1.0	9	14.9	1.0	9	14.2	1.0
05	Bl - Huckleberry - Thimbleberry	7	14.1	1.3		12.0		7	17.2	1.4
06	Bl - Oak fern - Heron's-bill	12	15.5	0.8	12	19.0	0.5	11	19.6	1.1
07	Bl - Devil's club - Lady fern		12.0			15.0			15.0	
08	Bl - Valerian - Sickle moss		9.0						9.0	
09	Bl - Horsetail - Glow moss		9.0						9.0	
10	Bl - Horsetail - Leafy moss		9.0					7	12.8	1.6

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] mk

Site Series(*)	Name	Ba			Bl			Hm			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	BlHm - Twistedstalk		12.0			12.0			12.0			12.0			12.0	
02	BlPa - Cladonia					9.0			9.0			9.0			9.0	
03	BlHm - Cladonia					9.0			9.0			9.0			9.0	
04	BlHm - Oak fern		12.0			12.0			12.0			12.0			12.0	
05	BlHm - Devil's club - Lady fern		12.0			12.0			12.0			15.0			15.0	
06	Bl - Horsetail - Leafy moss					9.0									9.0	
07	Bl - Lady fern - Horsetail					9.0									9.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] Sl mk

Site Series(*)	Name	Pa		
		#	Sl	SE
01	BlHm - Twistedstalk			
02	BlPa - Cladonia		9.0	
03	BlHm - Cladonia		9.0	
04	BlHm - Oak fern			
05	BlHm - Devil's club - Lady fern			
06	Bl - Horsetail - Leafy moss			
07	Bl - Lady fern - Horsetail			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g.,.18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, Sl = Mean Site Index, SE = Standard Error Of Mean

[redacted] mv3

Site Series(*)	Name	Bl			Pl			Se			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Bl - Rhododendron - Feathermoss		12.0			12.0			12.0				
02	BlPl - Crowberry - Cladina		9.0			9.0			9.0				
03	BlSb - Labrador tea		6.0			9.0			6.0			6.0	
04	Bl - Oak fern - Knight's plume		12.0			15.0			15.0				
05	Bl - Devil's club - Rhododendron		12.0			15.0			15.0				
06	Sxw - Huckleberry - Highbush-cranberry		12.0			12.0							
07	Bl - Horsetail - Feathermoss		9.0			9.0			9.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SI WV

Site Series(*)	Name	Bl			Pl			Se		
		#	SI	SE	#	SI	SE	#	SI	SE
01	BlHm - Azalea	9	11.9	1.2	7	15.1	0.8	10	14.9	1.7
02	BlPl - Cladonia					9.0			9.0	
03	BlHm - Feathermoss		9.0			9.0			9.0	
04	BlHm - Heron's-bill		12.0			12.0			12.0	
05	Bl - Oak fern - Heron's-bill		12.0			12.0			12.0	
06	Bl - Devil's club - Lady fern		12.0			15.0			15.0	
07	Bl - Valerian - Sickle moss		9.0						9.0	
08	Bl - Horsetail - Glow moss		9.0						9.0	
09	Bl - Lady fern - Horsetail		9.0						9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] 1

Site Series(*)	Name	Bl			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Hw - Step moss	8	18.1	0.8		15.0		7	20.8	0.8	7	20.7	1.4
02	Hw - Kinnikinnick - Cladonia		12.0			12.0			15.0				
03	HwBl - Oak fern		18.0			15.0			21.0		7	20.7	1.5
04	HwBl - Devil's club		18.0			18.0			24.0			24.0	
05	ActSx - Dogwood		21.0						21.0			24.0	
06	Hw - Azalea - Skunk cabbage		15.0			15.0						21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]1a

Site Series(*)	Name	Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Bramble		15.0			21.0			21.0	
02	HwBa - Oak fern		15.0			21.0			21.0	
03	HwBa - Devil's club - Lady fern		18.0			24.0			24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]2

Site Series(*)	Name	Bl			Cw			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Hw - Step moss	7	19.6	1.1	13	15.9	0.3	31	16.0	0.5	22	19.1	1.0	13	17.7	1.0
02	Hw - Kinnikinnick - Cladonia		12.0						12.0			15.0				
03	HwCw - Oak fern		18.0			15.0			15.0			21.0			21.0	
04	CwHw - Devil's club - Oak Fern		21.0			15.0			15.0			21.0			21.0	
05	Sx - Devil's club - Lady fern		21.0			18.0			18.0			24.0			24.0	
06	ActSx - Dogwood		21.0			21.0			18.0			21.0			24.0	
07	CwSx - Horsetail		15.0			18.0			15.0			15.0			21.0	
51	PlHw - Feathermoss								15.0			18.0				
52	SxEp - Thimbleberry- Hazelnut		18.0			15.0			15.0			21.0			21.0	
53	AtEp - Dogwood		18.0			15.0			15.0			21.0			21.0	
54	SxEp - Devil's club		21.0			18.0			15.0			21.0			24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Hw			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Hw - Devil's club		18.0			15.0			18.0	
02	Hw - Step moss		18.0			15.0			18.0	
03	Sx - Devil's club		18.0						18.0	
04	Sx - Devil's club - Dogwood		21.0						21.0	
05	ActSx - Dogwood		21.0						21.0	
06	Sx - Horsetail		15.0						18.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bl			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBl - Oak fern		18.0			15.0			21.0			21.0	
02	HwPl - Feathermoss - Cladonia					12.0			15.0				
03	Hw - Step moss		18.0			15.0			21.0			21.0	
04	HwBl - Devil's club		18.0			18.0			24.0			24.0	
05	Sx - Devil's club		18.0			18.0						24.0	
06	ActSx - Dogwood		21.0									24.0	
08	Sx - Horsetail		15.0			15.0						21.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Hm		
		#	SI	SE	#	SI	SE
01	HmBa - Blueberry		12.0			12.0	
02	HmBa - Mountain heather		8.0			8.0	
03	BaHm - Oak fern		12.0			12.0	
04	HmBa - Bramble		8.0			8.0	
05	BaHm - Twisted stalk		16.0			16.0	
06	HmYc - Deer cabbage		8.0			8.0	
07	YcHm - Hellebore		8.0			8.0	
08	HmYc - Sphagnum					8.0	
09	YcHm - Skunk cabbage					8.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Hm		
		#	SI	SE	#	SI	SE
01	HmBa - Blueberry		12.0			12.0	
02	HmBa - Mountain heather		8.0			8.0	
03	BaHm - Oak fern		12.0			12.0	
04	HmBa - Bramble		8.0			8.0	
05	BaHm - Twisted stalk		16.0			16.0	
06	HmYc - Deer cabbage		8.0			8.0	
07	YcHm - Hellebore		8.0			8.0	
08	HmYc - Sphagnum					8.0	
09	YcHm - Skunk cabbage					8.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] vh1

Site Series(*)	Name	Ba			Hm			Hw			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HmSs - Blueberry		12.0			12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]BPSmc

Site Series(*)	Name	Pl			Sb			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE
01	Pl - Feathermoss - Cladina		12.0			9.0			9.0	
02	Pl - Kinnikinnick - Cladonia		9.0						6.0	
03	SbPl - Feathermoss		9.0						6.0	
04	Sxw - Scrub birch - Feathermoss		9.0						6.0	
05	Sxw - Horsetail		15.0						12.0	
06	Sxw - Horsetail - Glow moss		12.0						9.0	
07	SbSxw - Scrub birch - Sedge		9.0						9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Pl			Sx			Sb			Fd		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Spirea - Purple peavine	24	19.5	0.4	17	18.0	0.6						
02	Pl - Juniper - Ricegrass		15.0			15.0							
03	Pl - Feathermoss - Cladina	13	16.2	0.6	10	14.9	0.6		9.0				
04	Fd - Soopolallie - Feathermoss		15.0			15.0							
05	Sxw - Spirea - Feathermoss	9	18.9	1.0		18.0							
06	Sxw - Twinberry - Coltsfoot	14	21.9	0.6	22	20.4	0.6					18.0	
07	Sxw - Horsetail		21.0			21.0							
08	Act - Dogwood - Prickly rose		21.0			21.0							
09	Sb - Creeping-snowberry - Sphagnum		12.0						9.0				
10	Sb - Soft-leaved sedge - Sphagnum		12.0			12.0			9.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bl			Pl			Sx			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Huckleberry	31	15.4	0.5	97	17.9	0.3	48	18.7	0.5			
02	Pl - Huckleberry - Cladonia		12.0		27	16.4	0.5	7	12.3	0.8			
03	SbPl - Feathermoss					15.0			15.0			9.0	
05	Sxw - Twinberry - Coltsfoot	11	16.3	1.0	26	19.1	0.5	35	21.3	0.6			
06	Sxw - Oak fern	17	19.1	0.7	10	19.9	0.5	26	19.3	0.5			
07	Sxw - Scrub birch - Feathermoss					12.0			12.0			9.0	
09	Sxw - Devil's club		18.0			21.0		8	19.8	0.8			
10	Sxw - Horsetail	8	15.3	0.9	7	19.6	0.8	9	19.3	1.0			
12	SbSxw - Scrub birch - Sedge					12.0			9.0			9.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] SBSwk3

Site Series(*)	Name	Bl			Pl			Sx			Fd			Sb		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Sxw - Oak fern	7	15.3	1.1	12	20.6	0.9	13	16.7	0.6						
02	Pl - Huckleberry - Cladina		12.0			15.0			12.0							
03	SxwFd - Purple peavine					15.0						15.0				
04	Sxw - Huckleberry - Highbush-cranberry		15.0			18.0			15.0							
05	Sb - Labrador tea				7	15.7	1.1		15.0					7	11.2	1.2
06	Sxw - Twinberry - Coltsfoot		15.0		7	20.7	0.8	7	19.2	1.4						
07	Sxw - Devil's club		18.0			21.0			21.0							
08	Sxw - Horsetail		18.0			21.0			18.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Bg			Cw			Fd			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Fd - Salal		20.0			20.0		12	29.7	2.3		20.0	
02	FdPl - Arbutus								16.0			16.0	
03	F[illegible] - Oniongrass								20.0			16.0	
04	[illegible]Bg - Oregon grape		24.0			24.0			28.0				
05	CwFd - Kindbergia					24.0			24.0				
06	CwBg - Foamflower		32.0			28.0			32.0				
07	Cw - Snowberry		32.0			28.0			32.0				
08	Act - Red-osier dogwood		24.0			24.0							
10	Pl - Sphagnum					8.0						16.0	
11	Cw - Skunk cabbage					16.0							
12	Cw - Vanilla leaf		28.0			24.0			28.0				
13	Cw - Indian plum		28.0			28.0			28.0				
14	Cw - Slough sedge		20.0			20.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Cw			Fd			Hw			Pl			Bg		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	Hw - Flat moss	18	27.6	0.6	12	34.1	1.2	13	30.3	0.7						
02	FdPl - Cladina					16.0						16.0				
03	FdHw - Salal		20.0			24.0										
04	Fd - Sword fern		24.0			28.0										
05	Cw - Sword Fern		28.0		11	39.7	0.7									
06	HwCw - Deer fern		32.0			36.0			32.0							
07	Cw - Foamflower	11	30.8	1.3	9	41.0	1.0	11	33.9	1.3					40.0	
08	Ss - Salmonberry		32.0												40.0	
09	Act - Red-osier dogwood		28.0												28.0	
11	Pl - Sphagnum		12.0									16.0				
12	CwSs - Skunk cabbage		20.0						32.0							
13	Cw - Salmonberry		28.0			36.0									36.0	
14	Cw - Black twinberry		28.0												28.0	
15	Cw - Slough sedge		24.0													

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ss		
		#	SI	SE
01	Hw - Flat moss			
02	FdPl - Cladina			
03	FdHw - Salal			
04	Fd - Sword fern			
05	Cw - Sword Fern			
06	HwCw - Deer fern			
07	Cw - Foamflower			
08	Ss - Salmonberry			
09	Act - Red-osier dogwood			
11	Pl - Sphagnum			
12	CwSs - Skunk cabbage		28.0	
13	Cw - Salmonberry			
14	Cw - Black twinberry			
15	Cw - Slough sedge			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Cw			Fd			Pl			Hw			Bg		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwFd - Cat's-tail moss		20.0		8	33.8	1.4									
02	FdPl - Kinnikinnick					12.0			12.0							
03	FdHw - Falsebox		12.0			16.0			16.0							
04	Fd - Fairybells		20.0			24.0										
05	Cw - Solomon's seal		24.0			36.0										
06	Hw - Queen's cup		24.0			28.0						28.0				
07	Cw - Devil's club		28.0			36.0									36.0	
08	Ss - Salmonberry		28.0												36.0	
09	Act - Red-osier dogwood		24.0												28.0	
11	Pl - Sphagnum		12.0						16.0							
12	CwSs - Skunk cabbage		20.0													

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Dr		
		#	SI	SE
01	HwFd - Cat's-tail moss			
02	FdPl - Kinnikinnick			
03	FdHw - Falsebox			
04	Fd - Fairybells			
05	Cw - Solomon's seal			
06	Hw - Queen's cup			
07	Cw - Devil's club	8	26.4	1.5
08	Ss - Salmonberry			
09	Act - Red-osier dogwood			
11	Pl - Sphagnum			
12	CwSs - Skunk cabbage			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Cw			Fd			Hw			Pl			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwFd - Cat's-tail moss		16.0			24.0			20.0							
02	FdPl - Kinnikinnick											12.0				
03	FdHw - Falsebox		12.0			16.0						16.0				
04	Fd - Fairybells		20.0			24.0						24.0				
05	Cw - Solomon's seal		24.0			36.0									32.0	
06	Hw - Queen's cup		24.0			28.0			28.0							
07	Cw - Devil's club		28.0			36.0									36.0	
08	Ss - Salmonberry		28.0												36.0	
09	Act - Red-osier dogwood		24.0												28.0	
11	Pl - Sphagnum		12.0									16.0				
12	CwSs - Skunk cabbage		20.0												24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Fd			Hw			Pl			Ba		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Pipecleaner moss		24.0			28.0			24.0							
02	FdHw - Salal					20.0						20.0				
03	HwCw - Salal		20.0			32.0										
04	CwHw - Sword fern		24.0			36.0										
05	BaCw - Foamflower		28.0			40.0			28.0							
06	HwBa - Deer fern		24.0						24.0						28.0	
07	BaCw - Salmonberry		28.0			36.0									28.0	
08	Ss - Salmonberry		28.0												28.0	
09	Act - Red-osier dogwood		28.0												24.0	
11	Pl - Sphagnum		12.0									16.0				
12	CwSs - Skunk cabbage		20.0						24.0							

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ss		
		#	SI	SE
01	HwBa - Pipecleaner moss			
02	FdHw - Salal			
03	HwCw - Salal			
04	CwHw - Sword fern			
05	BaCw - Foamflower			
06	HwBa - Deer fern			
07	BaCw - Salmonberry			
08	Ss - Salmonberry			
09	Act - Red-osier dogwood			
11	Pl - Sphagnum			
12	CwSs - Skunk cabbage		24.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] VHmm2

Site Series(*)	Name	Ba			Cw			Fd			Hw			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Pipecleaner moss		24.0			24.0			28.0			24.0			24.0	
02	FdHw - Salal								16.0							
03	HwCw - Salal					20.0			24.0						20.0	
04	CwHw - Sword fern					20.0			24.0						20.0	
05	BaCw - Foamflower		28.0			24.0			28.0			28.0			24.0	
06	HwBa - Deer fern		24.0			20.0						24.0			20.0	
07	CwYc - Goldthread					16.0									16.0	
08	BaCw - Salmonberry		28.0			24.0			28.0			24.0			24.0	
09	Pl - Sphagnum														12.0	
10	CwYc - Skunk cabbage					16.0						20.0			16.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pl		
		#	SI	SE
01	HwBa - Pipecleaner moss			
02	FdHw - Salal		16.0	
03	HwCw - Salal			
	CwHw - Sword fern			
0	BaCw - Foamflower			
06	HwBa - Deer fern			
07	CwYc - Goldthread			
08	BaCw - Salmonberry			
09	Pl - Sphagnum		16.0	
10	CwYc - Skunk cabbage			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Cw			Fd			Hw			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Step moss		16.0			20.0			24.0			20.0				
02	FdPl - Kinnikinnick								12.0						12.0	
03	FdHw - Falsebox								16.0							
04	BaCw - Oak fern		16.0			20.0			28.0			24.0				
05	HwBa - Queen's cup		20.0			20.0						24.0				
06	BaCw - Devil's club		24.0			24.0			28.0							
07	Ss - Salmonberry		24.0			24.0										
08	Act - Red-osier dogwood		20.0			20.0										
10	Pl - Sphagnum					12.0									12.0	
11	CwSs - Skunk cabbage					20.0										

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Yc			Bg		
		#	SI	SE	#	SI	SE
01	HwBa - Step moss						
02	FdPl - Kinnikinnick						
03	FdHw - Falsebox						
04	BaCw - Oak fern		20.0				
05	HwBa - Queen's cup		20.0				
06	BaCw - Devil's club		24.0			28.0	
07	Ss - Salmonberry						
08	Act - Red-osier dogwood						
10	Pl - Sphagnum						
11	CwSs - Skunk cabbage		20.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Cw			Fd			Hw			Pl			Ba		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Step moss		20.0			24.0			20.0							
02	FdPl - Kinnikinnick					12.0						12.0				
03	FdHw - Falsebox		16.0			16.0						16.0				
04	BaCw - Oak fern		20.0			28.0			20.0						16.0	
05	HwBa - Queen's cup		16.0						24.0						20.0	
06	BaCw - Devil's club		24.0			28.0									28.0	
07	Ss - Salmonberry		24.0												28.0	
08	Act - Red-osier dogwood		20.0												20.0	
10	Pl - Sphagnum		12.0									12.0				
11	CwSs - Skunk cabbage		20.0						20.0							

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted] /Whms2

Site Series(*)	Name	Ss		
		#	SI	SE
01	HwBa - Step moss			
02	FdPl - Kinnikinnick			
03	FdHw - Falsebox			
04	BaCw - Oak fern		24.0	
05	HwBa - Queen's cup			
06	BaCw - Devil's club		28.0	
07	Ss - Salmonberry		28.0	
08	Act - Red-osier dogwood			
10	Pl - Sphagnum			
11	CwSs - Skunk cabbage		20.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Cw			Hw			Pl			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Salal		12.0			16.0			16.0			16.0			20.0	
02	PlYc - Rhacomitrium					8.0						12.0				
03	CwYc - Salal					12.0			12.0			12.0				
04	HwSs - Lanky moss		24.0			20.0			24.0						24.0	
05	CwSs - Sword fern		28.0			24.0			28.0						28.0	
06	CwSs - Foamflower		24.0			24.0			24.0						32.0	
07	CwSs - Devil's club		28.0			24.0			28.0						32.0	
08	Ss - Lily-of-the-valley		28.0			24.0			28.0						28.0	
09	Ss - Trisetum		24.0			16.0			20.0						24.0	
11	CwYc - Goldthread					12.0			16.0							
12	PlYc - Sphagnum					8.0						8.0				
13	CwSs - Skunk cabbage					16.0			16.0						20.0	
14	Ss - Salal					8.0			12.0			12.0			12.0	
15	Ss - Kindbergia					16.0			20.0						24.0	
16	Ss - Reedgrass					12.0			20.0			20.0			24.0	
17	Ss - Sword fern					12.0			20.0						28.0	
18	Ss - Slough sedge					8.0									12.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Yc		
		#	SI	SE
01	CwHw - Salal			
02	PlYc - Rhacomitrium		8.0	
03	CwYc - Salal			
04	HwSs - Lanky moss			
05	CwSs - Sword fern			
06	CwSs - Foamflower			
07	CwSs - Devil's club			
08	Ss - Lily-of-the-valley			
09	Ss - Trisetum			
11	CwYc - Goldthread		8.0	
12	PlYc - Sphagnum		8.0	
13	CwSs - Skunk cabbage			
14	Ss - Salal			
15	Ss - Kindbergia			
16	Ss - Reedgrass			
17	Ss - Sword fern			
18	Ss - Slough sedge			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Cw			Hw			Pl			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	CwHw - Salal		16.0		7	19.7	0.8		20.0			20.0			20.0	
03	CwYc - Salal					16.0			16.0			16.0				
04	HwSs - Lanky moss	33	27.3	0.7	20	22.8	0.5	43	27.0	0.6				43	31.6	0.6
05	CwSs - Sword fern		24.0			24.0			24.0						28.0	
06	CwSs - Foamflower	15	31.0	0.7	10	24.2	0.5	18	28.2	0.7				27	32.9	0.8
07	CwSs - Devil's club		24.0		9	22.0	1.2	9	24.7	2.3				18	30.5	1.5
08	Ss - Lily-of-the-valley		24.0			24.0			24.0						32.0	
09	Ss - Trisetum								28.0						32.0	
11	CwYc - Goldthread					12.0			16.0			16.0				
12	PlYc - Sphagnum					8.0						12.0				
13	CwSs - Skunk cabbage					16.0			16.0						20.0	
14	Ss - Salal					12.0			12.0						16.0	
15	Ss - Kindbergia					16.0			16.0						24.0	
16	Ss - Reedgrass					12.0			12.0						16.0	
17	Ss - Sword fern					24.0			24.0						32.0	
18	Ss - Slough sedge					16.0									20.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Yc		
		#	SI	SE
01	CwHw - Salal		16.0	
03	CwYc - Salal			
04	HwSs - Lanky moss			
05	CwSs - Sword fern			
06	CwSs - Foamflower			
07	CwSs - Devil's club			
08	Ss - Lily-of-the-valley			
09	Ss - Trisetum			
11	CwYc - Goldthread		8.0	
12	PlYc - Sphagnum		8.0	
13	CwSs - Skunk cabbage		16.0	
14	Ss - Salal			
15	Ss - Kindbergia			
16	Ss - Reedgrass			
17	Ss - Sword fern			
18	Ss - Slough sedge			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Cw			Fd			Hw			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry	43	29.3	0.6	34	22.6	0.5	9	35.8	1.0	94	27.7	0.4	37	30.8	0.7
02	HwPl - Cladina					8.0			8.0			8.0				
03	HwCw - Salal					16.0		9	32.2	2.4		16.0				
04	CwHw - Sword fern		24.0		10	22.5	1.2		32.0		7	26.2	2.3		24.0	
05	BaCw - Foamflower	27	30.9	0.5	15	24.0	0.8		36.0		35	28.5	0.6	30	32.7	0.9
06	HwBa - Deer fern		24.0		9	23.3	1.9				20	25.2	1.4		24.0	
07	BaCw - Salmonberry		28.0			24.0					13	32.6	2.2		32.0	
08	BaSs - Devil's club		28.0		8	22.1	1.2				16	26.3	1.0	29	31.4	0.9
09	Ss - Salmonberry		28.0			24.0						28.0			28.0	
10	Act - Red-osier dogwood		28.0			24.0									32.0	
12	CwYc - Goldthread					12.0						16.0				
13	Pl - Sphagnum					8.0										
14	CwSs - Skunk cabbage				13	19.4	0.9				10	21.0	1.6	8	26.0	1.7

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Pl			Hm			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry									
02	HwPl - Cladina		12.0							
03	HwCw - Salal		16.0							
04	CwHw - Sword fern									
05	BaCw - Foamflower									
06	HwBa - Deer fern									
07	BaCw - Salmonberry									
08	BaSs - Devil's club									
09	Ss - Salmonberry									
10	Act - Red-osier dogwood									
12	CwYc - Goldthread		16.0			16.0			12.0	
13	Pl - Sphagnum		12.0							
14	CwSs - Skunk cabbage									

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

Site Series(*)	Name	Ba			Cw			Hw			Fd			Pl		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry		28.0			20.0			28.0							
02	HwPl - Cladina					8.0			8.0			8.0			12.0	
03	HwCw - Salal					16.0			16.0			24.0			20.0	
04	CwHw - Sword fern		20.0			20.0			24.0			24.0				
05	BaCw - Foamflower		28.0			24.0			28.0							
06	HwBa - Deer fern		24.0			20.0			24.0							
07	BaCw - Salmonberry		28.0			24.0			28.0							
08	BaSs - Devil's club		28.0			24.0			28.0							
09	CwYc - Goldthread					12.0			12.0						12.0	
10	Pl - Sphagnum														12.0	
11	CwYc - Skunk cabbage					16.0			16.0							

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Yc			Ss			Hm		
		#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Blueberry									
02	HwPl - Cladina		8.0							
03	HwCw - Salal		16.0							
04	CwHw - Sword fern		20.0			24.0				
05	BaCw - Foamflower		24.0			24.0				
06	HwBa - Deer fern		20.0			24.0				
07	BaCw - Salmonberry		24.0			28.0				
08	BaSs - Devil's club		24.0			28.0				
09	CwYc - Goldthread		12.0						12.0	
10	Pl - Sphagnum		8.0							
11	CwYc - Skunk cabbage		16.0			20.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Hw			Ss			Pl			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwSs - Lanky moss	15	21.4	0.6	11	22.9	0.9	36	29.2	1.1						
02	CwSs - Salal		20.0		7	23.9	1.1	11	27.9	1.8		20.0				
03	CwSs - Sword fern		20.0		8	26.9	0.9	31	29.7	0.9						
04	CwHw - Salal		16.0			24.0			24.0							
05	CwSs - Foamflower		20.0			28.0		32	35.4	0.6						
06	CwSs - Conocephalum		20.0			20.0		11	28.5	2.2						
07	Ss - Lily-of-the-valley		20.0			28.0			36.0							
08	Ss - Trisetum		12.0						32.0							
10	CwYc - Goldthread		12.0			16.0		8	31.6	2.0		16.0			16.0	
11	PlYc - Sphagnum		8.0									12.0			8.0	
12	CwSs - Skunk cabbage		12.0			16.0						16.0				
13	Ss - Salal		8.0			12.0			16.0			12.0				
14	Ss - Kindbergia		16.0			20.0			24.0							
15	Ss - Reedgrass		16.0			20.0			24.0			20.0				
16	Ss - Sword fern		20.0			28.0			36.0							
17	Ss - Slough sedge								36.0							

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Hw			Ss			Yc			Hm		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwSs - Lanky moss		16.0			20.0			24.0			16.0				
02	CwHw - Salal		16.0			16.0						16.0			12.0	
03	CwSs - Foamflower		16.0			20.0			24.0			16.0				
04	CwSs - Conocephalum		12.0			16.0			20.0			12.0				
05	CwYc - Goldthread		8.0			12.0						8.0			8.0	
06	CwSs - Skunk cabbage		8.0			12.0						8.0			8.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Cw			Hw			Pl			Sx		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwBa - Bramble		20.0			16.0			20.0			20.0			20.0	
02	Pl - Kinnikinnick					8.0			12.0			12.0				
03	HwPl - Feathermoss					16.0			16.0			16.0				
04	BaCw - Oak fern		24.0			16.0			24.0						24.0	
05	HwBa - Queen's cup		20.0			16.0			20.0						20.0	
06	BaCw - Devil's club		24.0			20.0			24.0						24.0	
07	Ss - Salmonberry		28.0			20.0			28.0						28.0	
08	Act - Red-osier dogwood		28.0			20.0									28.0	
10	Pl - Sphagnum					8.0						12.0				
11	CwSs - Skunk cabbage		12.0			12.0			12.0						12.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Fd		
		#	SI	SE
01	HwBa - Bramble			
02	Pl - Kinnikinnick			
03	HwPl - Feathermoss		20.0	
04	BaCw - Oak fern			
05	HwBa - Queen's cup			
06	BaCw - Devil's club			
07	Ss - Salmonberry			
08	Act - Red-osier dogwood			
10	Pl - Sphagnum			
11	CwSs - Skunk cabbage			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Fd			Hw			Pl			Bg		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HwFd - Kindbergia		24.0			32.0			24.0							
02	FdPl - Cladina					20.0						20.0				
03	FdHw - Salal		20.0			24.0						24.0				
04	Fd - Sword fern		28.0			28.0										
05	Cw - Sword fern		24.0			36.0									32.0	
06	HwCw - Deer fern		24.0			32.0			24.0							
07	Cw - Foamflower		28.0			40.0									40.0	
08	Ss - Salmonberry		24.0												40.0	
09	Act - Red-osier dogwood		24.0												28.0	
11	Pl - Sphagnum		8.0									16.0				
12	CwSs - Skunk cabbage		16.0						32.0							
13	Cw - Salmonberry		24.0			36.0									36.0	
14	Cw - Black twinberry		28.0												28.0	
15	Cw - Slough sedge		24.0													

continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ss		
		#	SI	SE
01	HwFd - Kindbergia			
02	FdPl - Cladina			
03	FdHw - Salal			
04	Fd - Sword fern			
05	Cw - Sword fern			
06	HwCw - Deer fern			
07	Cw - Foamflower			
08	Ss - Salmonberry			
09	Act - Red-osier dogwood			
11	Pl - Sphagnum			
12	CwSs - Skunk cabbage		28.0	
13	Cw - Salmonberry			
14	Cw - Black twinberry			
15	Cw - Slough sedge			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Bl			Pl			Se			Fd		
		#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE	#	Sl	SE
01	BlBa - Rhododendron		15.0			15.0			15.0			15.0				
02	BlPl - Juniper - Rhacomitrium								12.0			12.0				
03	FdBl - Falsebox - Pinegrass								15.0			12.0			15.0	
04	Bl - Huckleberry - Falsebox					12.0			15.0			12.0			15.0	
05	BlBa - Azalea - Pipecleaner moss		15.0			15.0			18.0			15.0				
06	Bl - Gooseberry - Valerian		15.0			15.0			15.0			15.0				
07	BlBa - Oak fern - Lady fern		18.0			18.0						18.0				
08	Bl - Gooseberry - Horsetail					9.0						18.0				

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, Sl = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Cw			Fd			Pl			Py			Ss		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	FdCw - Hazelnut		15.0			24.0			24.0			24.0				
02	FdPl - Peltigera					12.0			12.0							
03	Fd - Falsebox - Feathermoss					18.0			18.0			18.0				
04	Fd - Douglas maple - Fairybells		15.0			24.0						24.0				
05	CwFd - Vine maple		24.0			27.0									27.0	
06	Cw - Devil's club - Lady fern		27.0			36.0									36.0	
07	CwSxw - Skunk cabbage		15.0												21.0	

...continued

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Bg		
		#	SI	SE
01	FdCw - Hazelnut			
02	FdPl - Peltigera			
03	Fd - Falsebox - Feathermoss			
04	Fd - Douglas maple - Fairybells			
05	CwFd - Vine maple			
06	Cw - Devil's club - Lady fern		36.0	
07	CwSxw - Skunk cabbage			

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Hm		
		#	SI	SE	#	SI	SE
01	HmBa - Blueberry		12.0			16.0	
02	HmBa - Mountain heather		8.0			12.0	
03	BaHm - Oak fern		12.0				
04	HmBa - Bramble		12.0			16.0	
05	BaHm - Twisted stalk		16.0			16.0	
06	HmYc - Deer cabbage		8.0			8.0	
07	YcHm - Hellebore		8.0			8.0	
08	HmYc - Sphagnum					8.0	
09	YcHm - Skunk cabbage					8.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Hm		
		#	SI	SE	#	SI	SE
01	HmBa - Blueberry		16.0			16.0	
02	HmBa - Mountain heather		12.0			12.0	
03	BaHm - Oak fern		16.0			16.0	
04	HmBa - Bramble		16.0			16.0	
05	BaHm - Twisted stalk		16.0			16.0	
06	HmYc - Deer cabbage		8.0			8.0	
07	YcHm - Hellebore		8.0			8.0	
08	HmYc - Sphagnum					8.0	
09	YcHm - Skunk cabbage					8.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:
= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean

[redacted]

Site Series(*)	Name	Ba			Hm			Hw			Yc		
		#	SI	SE	#	SI	SE	#	SI	SE	#	SI	SE
01	HmSs - Blueberry		12.0			12.0			12.0			12.0	

* Restrictions apply to some species. See the appropriate Establishment to Free Growing guidebook, May, 2000.

** The SIBEC reports contain two types of site index estimates for each site series/tree species combination: a single class value where the first approximation site index class (BC MoF 1997) is reported, e.g., 18.0 or three values - sample size, mean site index, and standard error - where an updated post-1997 estimate is presented.

Legend of Terms:

= Sample Size, SI = Mean Site Index, SE = Standard Error Of Mean